

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/22, 16/12, C12N 15/70, A61K 38/16, 39/095, G01N 33/569, C12Q 1/68, C12N 1/21 // (C12N 1/21, C12R 1:19)	A1	(11) International Publication Number: WO 96/29412 (43) International Publication Date: 26 September 1996 (26.09.96)
(21) International Application Number: PCT/CA96/00157 (22) International Filing Date: 15 March 1996 (15.03.96) (30) Priority Data: 08/406,362 17 March 1995 (17.03.95) US 60/001,983 4 August 1995 (04.08.95) US (71) Applicant (for all designated States except US): IAF BIO VAC INC. [CA/CA]; 525 Des Prairies Boulevard, Laval, Quebec H7N 4Z2 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): BRODEUR, Bernard, R. [CA/CA]; 2401 Maritain Street, Sillery, Quebec G1T 1N6 (CA). MARTIN, Denis [CA/CA]; 4728-G Gaboury Street, St-Augustin-de-Des-Maures, Quebec G34 1E9 (CA). HAMEL, Josée [CA/CA]; 2401 Maritain Street, Sillery, Quebec G1T 1N6 (CA). RIOUX, Clément [CA/CA]; 1012 Jean Charles Cantin, Ville-de-Cap-Rouge, Quebec G1Y 2X1 (CA).		(74) Agents: DUBUC, Jean et al.; Goudreau Gage Dubuc & Martineau Walker, 800 The Stock Exchange Tower, Victoria Square, P.O. Box 242, Montreal, Quebec H4Z 1E9 (CA). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROTEINASE K RESISTANT SURFACE PROTEIN OF NEISSERIA MENINGITIDIS (57) Abstract A highly conserved, immunologically accessible antigen at the surface of <i>Neisseria meningitidis</i> organisms. Immunotherapeutic, prophylactic and diagnostic compositions and methods useful in the treatment, prevention and diagnosis of <i>Neisseria meningitidis</i> diseases. A proteinase K resistant <i>Neisseria meningitidis</i> surface protein having an apparent molecular weight of 22 kDa, the corresponding nucleotide and derived amino acid sequences (SEQ ID NO:1, NO:3, NO:5, and NO:7: SEQ ID NO:2, NO:4, NO:6, and NO:8), recombinant DNA methods for the production of the <i>Neisseria meningitidis</i> 22 kDa surface protein, and antibodies that bind to the <i>Neisseria meningitidis</i> 22 kDa surface protein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**PROTEINASE K RESISTANT SURFACE
PROTEIN OF *NEISSERIA MENINGITIDIS***

5 TECHNICAL FIELD OF THE INVENTION

This invention relates to a highly conserved, immunologically accessible antigen at the surface of *Neisseria meningitidis* organisms. This unique antigen provides the basis for new immunotherapeutic, prophylactic and diagnostic agents useful in the treatment, prevention and diagnosis of *Neisseria meningitidis* diseases. More particularly, this invention relates to a proteinase K resistant *Neisseria meningitidis* surface protein having an apparent molecular weight of 22 kDa, the corresponding nucleotide and derived amino acid sequences (SEQ ID NO:1 to SEQ ID NO:26), recombinant DNA methods for the production of the *Neisseria meningitidis* 22 kDa surface protein, antibodies that bind to the *Neisseria meningitidis* 22 kDa surface protein and methods and compositions for the diagnosis, treatment and prevention of *Neisseria meningitidis* diseases.

BACKGROUND OF THE INVENTION

Neisseria meningitidis is a major cause of death and morbidity throughout the world. *Neisseria meningitidis* causes both endemic and epidemic diseases, principally meningitis and meningococemia [Gold, Evolution of meningococcal disease, p. 69, Vedros N.A., CRC Press (1987); Schwartz et al., Clin. Microbiol. Rev., 2, p. S118 (1989)]. In fact, this organism is one of the most common causes, after *Haemophilus influenzae* type b, of bacterial meningitis in the United States, accounting for approximately 20% of all cases. It has been well documented that serum bactericidal activity is the major defense mechanism against *Neisseria meningitidis* and that

protection against invasion by the bacteria correlates with the presence in the serum of anti-meningococcal antibodies [Goldschneider et al., J. Exp. Med. 129, p. 1307 (1969); Goldschneider et al., J. Exp. Med., 129, 5 p. 1327 (1969)].

Neisseria meningitidis are subdivided into serological groups according to the presence of capsular antigens. Currently, 12 serogroups are recognized, but serogroups A, B, C, Y, and W-135 are most commonly found. 10 Within serogroups, serotypes, subtypes and immunotypes can be identified on outer membrane proteins and lipopolysaccharide [Frasch et al., Rev. Infect. Dis. 7, p. 504 (1985)].

The capsular polysaccharide vaccines presently 15 available are not effective against all *Neisseria meningitidis* isolates and do not effectively induce the production of protective antibodies in young infants [Frasch, Clin. Microbiol. Rev. 2, p. S134 (1989); Reingold et al., Lancet, p. 114 (1985); Zollinger, in Woodrow and 20 Levine, New generation vaccines, p. 325, Marcel Dekker Inc. N.Y. (1990)]. The capsular polysaccharide of serogroups A, C, Y and W-135 are presently used in vaccines against this organism. These polysaccharide vaccines are effective in the short term, however the 25 vaccinated subjects do not develop an immunological memory, so they must be revaccinated within a three-year period to maintain their level of resistance.

Furthermore, these polysaccharide vaccines do not induce sufficient levels of bactericidal antibodies to 30 obtain the desired protection in children under two years of age, who are the principal victims of this disease. No effective vaccine against serogroup B isolates is presently available even though these organisms are one of the primary causes of meningococcal diseases in developed 35 countries. Indeed, the serogroup B polysaccharide is not

a good immunogen, inducing only a poor response of IgM of low specificity which is not protective [Gotschlich et al., J. Exp. Med., p. 129, 1349 (1969); Skevakis et al., J. Infect. Dis., 149, p. 387 (1984); Zollinger et al., J. Clin. Invest., 63, p. 836 (1979)]. Furthermore, the presence of closely similar, crossreactive structures in the glycoproteins of neonatal human brain tissue [Finne et al., Lancet, p. 355 (1983)] might discourage attempts at improving the immunogenicity of serogroup B polysaccharide.

To obtain a more effective vaccine, other *Neisseria meningitidis* surface antigens such as lipopolysaccharide, pili proteins and proteins present in the outer membrane are under investigation. The presence of a human immune response and bactericidal antibodies against certain of these proteinaceous surface antigens in the sera of immunized volunteers and convalescent patients was demonstrated [Mandrell and Zollinger, Infect. Immun., 57, p. 1590 (1989); Poolman et al., Infect. Immun., 40, p. 398 (1983); Rosenquist et al., J. Clin. Microbiol., 26, p. 1543 (1988); Wedege and Froholm, Infect. Immun., 51, p. 571 (1986); Wedege and Michaelsen, J. Clin. Microbiol., 25, p. 1349 (1987)].

Furthermore, monoclonal antibodies directed against outer membrane proteins, such as class 1, 2/3 and 5, were also reported to be bactericidal and to protect against experimental infections in animals [Brodeur et al., Infect. Immun., 50, p. 510 (1985); Frasch et al., Clin. Invest. Med., 9, p. 101 (1986); Saukkonen et al., Microb. Pathogen., 3, p. 261 (1987); Saukkonen et al., Vaccine, 7, p. 325 (1989)].

Antigen preparations based on *Neisseria meningitidis* outer membrane proteins have demonstrated immunogenic effects in animals and in humans and some of them have been tested in clinical trials [Bjune et al., Lancet,

p. 1093 (1991); Costa et al., NIPH Annals, 14, p. 215 (1991); Frascch et al., Med. Trop., 43, p. 177 (1982); Frascch et al., Eur. J. Clin. Microbiol., 4, p. 533 (1985); Frascch et al. in Robbins, Bacterial Vaccines, p. 262, Praeger Publications, N.Y. (1987); Frascch et al., J. Infect. Dis., 158, p. 710 (1988); Moreno et al., Infect. Immun., 47, p. 527 (1985); Rosenqvist et al., J. Clin. Microbiol., 26, p. 1543 (1988); Sierra et al., NIPH Annals, 14, p. 195 (1991); Wedege and Froholm, Infect. Immun., 51, p. 571 (1986); Wedege and Michaelsen, J. Clin. Microbiol., 25, p. 1349 (1987); Zollinger et al., J. Clin. Invest., 63, p. 836 (1979); Zollinger et al., NIPH Annals, 14, p. 211 (1991)]. However, the existence of great interstrain antigenic variability in the outer membrane proteins can limit their use in vaccines [Frascch, Clin. Microb., Rev. 2, p. S134 (1989)]. Indeed, most of these preparations induced bactericidal antibodies that were restricted to the same or related serotype from which the antigen was extracted [Zollinger in Woodrow and Levine, New Generation Vaccines, p. 325, Marcel Dekker Inc. N.Y. (1990)]. Furthermore, the protection conferred by these vaccines in young children has yet to be clearly established. The highly conserved *Neisseria meningitidis* outer membrane proteins such as the class 4 [Munkley et al., Microb. Pathogen., 11, p. 447 (1991)] and the lip protein (also called H.8) [Woods et al., Infect. Immun., 55, p. 1927 (1987)] are not interesting vaccine candidates since they do not elicit the production of bactericidal antibodies. To improve these vaccine preparations, there is a need for highly conserved proteins that would be present at the surface of all *Neisseria meningitidis* strains and that would be capable of eliciting bactericidal antibodies in order to develop a broad spectrum vaccine.

The current laboratory diagnosis of *Neisseria meningitidis* is usually done by techniques such as Gram stain of smear preparations, latex agglutination or coagglutination, and the culture and isolation on enriched and selective media [Morello et al., in Balows, Manual of Clinical Microbiology, p. 258, American Society for Microbiology, Washington (1991)]. Carbohydrate degradation tests are usually performed to confirm the identification of *Neisseria meningitidis* isolates. Most of the described procedures are time-consuming processes requiring trained personnel. Commercial agglutination or coagglutination kits containing polyvalent sera directed against the capsular antigens expressed by the most prevalent serogroups are used for the rapid identification of *Neisseria meningitidis*. However, these polyvalent sera often nonspecifically cross-react with other bacterial species and for that reason should always be used in conjunction with Gram stain and culture. Accordingly, there is a need for efficient alternatives to these diagnostic assays that will improve the rapidity and reliability of the diagnosis.

DISCLOSURE OF THE INVENTION

The present invention solves the problems referred to above by providing a highly conserved, immunologically accessible antigen at the surface of *Neisseria meningitidis* organisms. Also provided are recombinant DNA molecules that code for the foregoing antigen, unicellular hosts transformed with those DNA molecules, and a process for making substantially pure, recombinant antigen. Also provided are antibodies specific to the foregoing *Neisseria meningitidis* antigen. The antigen and antibodies of this invention provide the basis for unique methods and pharmaceutical compositions for the detection,

prevention and treatment of *Neisseria meningitidis* diseases.

The preferred antigen is the *Neisseria meningitidis* 22 kDa surface protein, including fragments, analogues and derivatives thereof. The preferred antibodies are the Me-1 and Me-7 monoclonal antibodies specific to the *Neisseria meningitidis* 22 kDa surface protein. These antibodies are highly bacteriolytic against *Neisseria meningitidis* and passively protect mice against experimental infection.

10 The present invention further provides methods for isolating novel *Neisseria meningitidis* surface antigens and antibodies specific thereto.

BRIEF DESCRIPTION OF THE DRAWINGS

15 Figure 1 depicts the nucleotide and derived amino acid sequences of the *Neisseria meningitidis* strain 608B 22 kDa surface protein (SEQ ID NO:1; SEQ ID NO:2). Conventional three letter symbols are used for the amino acid residues. The open reading frame extends from the start codon at base 143 to the stop codon at base 667. The box indicates the putative ribosome binding site whereas the putative -10 promoter sequence is underlined. A 19-amino-acid signal peptide is also underlined.

25 Figure 2 is a photograph of a Coomassie Blue stained 14% SDS-PAGE gel displaying α -chymotrypsin and trypsin digests of *Neisseria meningitidis* strain 608B (B:2a:P1.2) outer membrane preparations. Lane 1 contains the following molecular weight markers: Phosphorylase b (97,400); bovine serum albumin (66,200); ovalbumin (45,000); carbonic anhydrase (31,000); soybean trypsin inhibitor (21,500); and lysozyme (14,400). Lane 2 contains undigested control outer membrane preparation. Lane 3 contains α -chymotrypsin treated preparation (2mg of

enzyme per mg of protein); lane 4 contains trypsin treated preparation.

Figure 3a is a photograph of a Coomassie Blue stained 14% SDS-PAGE gel displaying proteinase K digests of *Neisseria meningitidis* strain 608B (B:2a:P1.2) outer membrane preparations. Lanes 1, 3, 5, and 7 contain undigested control. Lanes 2, 4, 6 and 8 contain outer membrane preparations digested with proteinase K (2 IU per mg of protein). Lanes 1 to 4 contain preparations treated at pH 7.2. Lanes 5 to 8 contain preparations treated at pH 9.0. Lanes 1, 2, 5 and 6 contain preparations treated without SDS. Lanes 3, 4, 7 and 8 contain preparations treated in the presence of SDS. Molecular weight markers are indicated on the left (in kilodaltons).

Figure 3b is a photograph of a Coomassie Blue stained 14% SDS-PAGE gel displaying the migration profiles of affinity purified recombinant 22 kDa protein. Lane 1 contains molecular weight markers: Phosphorylase b (97,400), bovine serum albumin (66,200), ovalbumin (45,000), carbonic anhydrase (31,000), soybean trypsin inhibitor (21,500) and lysozyme (14,400). Lane 2 contains 5 µg of control affinity purified recombinant 22 kDa protein. Lane 3 contains 5 µg of affinity purified recombinant 22 kDa protein heated at 100°C for 5 min. . Lane 4 contains 5 µg of affinity purified recombinant 22 kDa protein heated at 100°C for 10 min. . Lane 5 contains 5 µg of affinity purified recombinant 22 kDa protein heated at 100°C for 15 min.

Figure 4 is a photograph of Coomassie Blue stained 14% SDS-PAGE gels and their corresponding Western immunoblots showing the reactivity of monoclonal antibodies specific to the *Neisseria meningitidis* 22 kDa surface protein. Outer membrane preparation from *Neisseria meningitidis* strain 608B (B:2a:P1.2) (A) untreated; (B) Proteinase K treated (2 IU per mg of

protein). Lane 1 contains molecular weight markers and characteristic migration profile on 14% SDS-PAGE gel of outer membrane preparations. Lane 2 contains Me-2; Lane 3 contains Me-3; lane 4 contains Me-5; lane 5 contains Me-7; and lane 6 contains an unrelated control monoclonal antibody. The molecular weight markers are phosphorylase b (97,400), bovine serum albumin (66,200), ovalbumin (45,000), carbonic anhydrase (31,000), soybean trypsin inhibitor (21,500) and lysozyme (14,400). The immunoblot results shown in Figure 4 for Me-2, Me-3, Me-5, Me-6 and Me-7 are consistent with the immunoblot results obtained for Me-1.

Figure 5 is a graphic depiction of the binding activity of the monoclonal antibodies to intact bacterial cells. The results for representative monoclonal antibodies Me-5 and Me-7 are presented in counts per minute ("CPM") on the vertical axis. The various bacterial strains used in the assay are shown on the horizontal axis. A *Haemophilus influenzae* porin-specific monoclonal antibody was used as a negative control. Background counts below 500 CPM were recorded and were subtracted from the binding values.

Figure 6 is a photograph of stained 14% SDS-PAGE gels and their corresponding Western immunoblot demonstrating the purification of the recombinant 22 kDa *Neisseria meningitidis* surface protein from concentrated culture supernatant of *Escherichia coli* strain BL21(DE3). Figure 6(A) is a photograph of a Coomassie Blue and silver stained 14% SDS-Page gel. Lane 1 contains the following molecular weight markers: phosphorylase b (97,400), bovine serum albumin (66,200), ovalbumin (45,000), carbonic anhydrase (31,000), soybean trypsin inhibitor (21,500) and lysozyme (14,400). Lane 2 contains outer membrane protein preparation extracted from *Neisseria meningitidis* strain 608B (serotype B:2a:p1.2) (10 mg). Lane 3 contains

concentrated culture supernatant of *Escherichia coli* BL21(DE3) (10 mg). Lane 4 contains affinity purified recombinant 22 kDa *Neisseria meningitidis* surface protein (1 mg). Figure 6(B) is a photograph of a Coomassie Blue stained 14% SDS-PAGE gel of α -chymotrypsin, trypsin and proteinase K digests of affinity purified recombinant 22 kDa *Neisseria meningitidis* surface protein. Lane 1 contains the following molecular weight markers: phosphorylase b (97,400), bovine serum albumin (66,200), ovalbumin (45,000), carbonic anhydrase (31,000), soybean trypsin inhibitor (21,500) and lysozyme (14,400). Lanes 2 to 5 contain purified recombinant 22 kDa *Neisseria meningitidis* surface protein (2 mg). Lanes 7 to 10 contain bovine serum albumin (2 mg). Lanes 2 and 7 contain undigested protein ("NT"). Lanes 3 and 8 contain α -chymotrypsin ("C") treated protein (2 mg of enzyme per mg of protein). Lanes 4 and 9 contain trypsin ("T") treated protein (2 mg of enzyme per mg of protein). Lanes 5 and 10 contain proteinase K ("K") treated protein (2 IU per mg of protein). Figure 6(C) is a photograph of the Western immunoblotting of a duplicate gel using monoclonal antibody Me-5.

Figure 7 is a graphical depiction of the bactericidal activity of protein A-purified anti-*Neisseria meningitidis* 22 kDa surface protein monoclonal antibodies against *Neisseria meningitidis* strain 608B (B:2a:P1.2). The vertical axis of the graph shows the percentage of survival of the *Neisseria meningitidis* bacteria after exposure to various concentrations of monoclonal antibody (shown on the horizontal axis of the graph).

Figure 8 depicts the nucleotide and derived amino acid sequences of the *Neisseria meningitidis* strain MCH88 22 kDa surface protein (SEQ ID NO:3; SEQ ID NO:4). Conventional three letter symbols are used for the amino

acid residues. The open reading frame extends from the start codon at base 116 to the stop codon at base 643.

Figure 9 depicts the nucleotide and derived amino acid sequences of the *Neisseria meningitidis* strain Z4063 22 kDa surface protein (SEQ ID NO:5; SEQ ID NO:6). Conventional three letter symbols are used for the amino acid residues. The open reading frame extends from the start codon at base 208 to the stop codon at base 732.

Figure 10 depicts the nucleotide and derived amino acid sequences of the *Neisseria gonorrhoeae* strain b2, 22 kDa surface protein (SEQ ID NO:7; SEQ ID NO:8). Conventional three letter symbols are used for the amino acid residues. The open reading frame extends from the start codon at base 241 to the stop codon at base 765.

Figure 11 depicts the consensus sequence established from the DNA sequences of the four strains of *Neisseria* and indicates the substitutions or insertion of nucleotides specific to each strain.

Figure 12 depicts the consensus sequence established from the protein sequences of the four strains of *Neisseria* and indicates the substitutions or insertion of amino acid residues specific to each strain.

Figure 13 represents the time course of the immune response to the recombinant 22 kDa protein in rabbits expressed as the reciprocal ELISA titer. The rabbits were injected with outer membrane preparations from *E. coli* strain JM109 with plasmid pN2202 or with control plasmid pWKS30. The development of the specific humoral response was analysed by ELISA using outer membrane preparations obtained from *Neisseria meningitidis* strain 608B (B:2a:P1.2) as coating antigen.

Figure 14 represents the time course of the immune response to the recombinant 22kDa protein in *Macaca fascicularis* (cynomolgus) monkeys expressed as the reciprocal ELISA titer. The two monkeys were respectively

immunized with 100µg (K28) and 200µg (I276) of affinity purified 22kDa protein per injection. The control monkey (K65) was immunized with 150µg of unrelated recombinant protein following the same procedure. The development of
5 the specific humoral response was analysed by ELISA using outer membrane preparations obtained from *Neisseria meningitidis* strain 608B (B:2a:P1.2) as coating antigen.

Figure 15 is a graphic representation of the synthetic peptides of the invention as well as their
10 respective position in the full 22kDa protein of *Neisseria meningitidis* strain 608B (B:2a:P1.2).

Figure 16 is a map of plasmid pNP2204 containing the gene which encodes the *Neisseria meningitidis* 22 kDa surface protein 22kDa, *Neisseria meningitidis* 22 kDa
15 surface protein gene; Amp^r, ampicillin-resistance coding region; ColE1, origin of replication; cI857, bacteriophage λ cI857 temperature-sensitive repressor gene; λPL, bacteriophage λ transcription promoter; T1 transcription terminator. The direction of transcription is indicated by
20 the arrows. BglIII and BamHI are the restriction sites used to insert the 22 kDa gene in the p629 plasmid.

DETAILED DESCRIPTION OF THE INVENTION

During our study of the ultrastructure of the outer
25 membrane of *Neisseria meningitidis* we identified a new low molecular weight protein of 22 kilodaltons which has very unique properties. This outer membrane protein is highly resistant to extensive treatments with proteolytic enzymes, such as proteinase K, a serine protease derived
30 from the mold *Tritirachium album* limber. This is very surprising since proteinase K resistant proteins are very rare in nature because of the potency, wide pH optimum, and low peptide bond specificity of this enzyme [Barrett, A.J. and N.D. Rawlings, Biochem. Soc. Transactions (1991)]

19: 707-715]. Only a few reports have described proteins of prokaryotic origin that are resistant to the enzymatic degradation of proteinase K. Proteinase K resistant proteins have been found in *Leptospira* species [Nicholson, 5 V.M. and J.F. Prescott, Veterinary Microbiol. (1993) 36:123-138], *Mycoplasma* species [Butler, G.H. et al. Infec. Immun. (1991) 59:1037-1042], *Spiroplasma mirum* [Bastian, F.O. et al. J. Clin. Microbiol. (1987) 25:2430-2431] and in viruses and prions [Onodera, T. et al. 10 Microbiol. Immunol. (1993) 37:311-316; Prusiner, S.B. et al. Proc. Nat. Acad. Sci. USA (1993) 90:2793-2797]. Herein, we describe the use of this protein as a means for the improved prevention, treatment and diagnosis of *Neisseria meningitidis* infections.

15 Thus according to one aspect of the invention we provide a highly conserved, immunologically accessible *Neisseria meningitidis* surface protein, and fragments, analogues, and derivatives thereof. As used herein, "Neisseria meningitidis surface protein" means any 20 *Neisseria meningitidis* surface protein encoded by a naturally occurring *Neisseria meningitidis* gene. The *Neisseria meningitidis* protein according to the invention may be of natural origin, or may be obtained through the application of molecular biology with the object of 25 producing a recombinant protein, or fragment thereof.

As used herein, "highly conserved" means that the gene for the *Neisseria meningitidis* surface protein and the protein itself exist in greater than 50% of known strains of *Neisseria meningitidis*. Preferably, the gene 30 and its protein exist in greater than 99% of known strains of *Neisseria meningitidis*. Examples 2 and 4 set forth methods by which one of skill in the art would be able to test a variety of different *Neisseria meningitidis* surface proteins to determine if they are "highly conserved".

As used herein, "immunologically accessible" means that the *Neisseria meningitidis* surface protein is present at the surface of the organism and is accessible to antibodies. Example 2 sets forth methods by which one of skill in the art would be able to test a variety of different *Neisseria meningitidis* surface proteins to determine if they are "immunologically accessible". Immunological accessibility may be determined by other methods, including an agglutination assay, an ELISA, a RIA, an immunoblotting assay, a dot-enzyme assay, a surface accessibility assay, electron microscopy, or a combination of these assays.

As used herein, "fragments" of the *Neisseria meningitidis* surface protein include polypeptides having at least one peptide epitope, or analogues and derivatives thereof. Peptides of this type may be obtained through the application of molecular biology or synthesized using conventional liquid or solid phase peptide synthesis techniques.

As used herein, "analogues" of the *Neisseria meningitidis* surface protein include those proteins, or fragments thereof, wherein one or more amino acid residues in the naturally occurring sequence is replaced by another amino acid residue, providing that the overall functionality and protective properties of this protein are preserved. Such analogues may be produced synthetically, or by recombinant DNA technology, for example, by mutagenesis of a naturally occurring *Neisseria meningitidis* surface protein. Such procedures are well known in the art.

For example, one such analogue is selected from the recombinant protein that may be produced from the gene for the 22kDa protein from *Neisseria gonorrhoeae* strain b2, as depicted in Figure 10. A further analog may be obtained

from the isolation of the corresponding gene from *Neisseria lactamica*.

As used herein, a "derivative" of the *Neisseria meningitidis* surface protein is a protein or fragment thereof that has been covalently modified, for example, with dinitrophenol, in order to render it immunogenic in humans. The derivatives of this invention also include derivatives of the amino acid analogues of this invention.

It will be understood that by following the examples of this invention, one of skill in the art may determine without undue experimentation whether a particular fragment, analogue or derivative would be useful in the diagnosis, prevention or treatment of *Neisseria meningitidis* diseases.

This invention also includes polymeric forms of the *Neisseria meningitidis* surface proteins, fragments, analogues and derivatives. These polymeric forms include, for example, one or more polypeptides that have been crosslinked with crosslinkers such as avidin/biotin, glutaraldehyde or dimethylsuberimide. Such polymeric forms also include polypeptides containing two or more tandem or inverted contiguous *Neisseria meningitidis* sequences, produced from multicistronic mRNAs generated by recombinant DNA technology.

This invention provides substantially pure *Neisseria meningitidis* surface proteins. The term "substantially pure" means that the *Neisseria meningitidis* surface protein according to the invention is free from other proteins of *Neisseria meningitidis* origin. Substantially pure *Neisseria meningitidis* surface protein preparations can be obtained by a variety of conventional processes, for example the procedure described in Examples 3 and 11.

In a further aspect, the invention particularly provides a 22 kDa surface protein of *Neisseria meningitidis* having the amino acid sequence of Figure 1

(SEQ ID NO:2), or a fragment, analogue or derivative thereof.

In a further aspect, the invention particularly provides a 22 kDa surface protein of *Neisseria*

5 *meningitidis* having the amino acid sequence of Figure 8 (SEQ ID NO:4), Figure 9 (SEQ ID NO:6) or a fragment, analogue or derivative thereof. Such a fragment may be selected from the peptides listed in Figure 15 (SEQ ID NO:9 to SEQ ID NO:26).

10 In a further aspect, the invention provides a 22kDa surface protein of *Neisseria gonorrhoeae* having the amino acid sequence of Figure 10 (SEQ ID NO:8), or a fragment, analogue or derivative thereof. As will be apparent from the above, any reference to the *Neisseria meningitidis*
15 22kDa protein also encompasses 22kDa proteins isolated from, or made from genes isolated from other species of *Neisseriaceae* such as *Neisseria gonorrhoeae* or *Neisseria lactamica*.

A *Neisseria meningitidis* 22 kDa surface protein
20 according to the invention may be further characterized by one or more of the following features:

(1) it has an approximate molecular weight of 22 kDa as evaluated on SDS-PAGE gel;

(2) its electrophoretic mobility on SDS-PAGE gel is
25 not modified by treatment with reducing agents;

(3) it has an isoelectric point (pI) in a range around pI 8 to pI 10;

(4) it is highly resistant to degradation by proteolytic enzymes such as α -chymotrypsin, trypsin and
30 proteinase K;

(5) periodate oxidation does not abolish the specific binding of antibody directed against the *Neisseria meningitidis* 22 kDa surface protein;

(6) it is a highly conserved antigen;

(7) it is accessible to antibody at the surface of intact *Neisseria meningitidis* organisms;

(8) it can induce the production of bactericidal antibodies;

5 (9) it can induce the production of antibodies that can protect against experimental infection;

(10) it can induce, when injected into an animal host, the development of an immunological response that can protect against *Neisseria meningitidis* infection.

10 This invention also provides, for the first time, a DNA sequence coding for the *Neisseria meningitidis* 22 kDa surface protein (SEQ ID NO:1, NO:3, NO:5, and NO:7). The preferred DNA sequences of this invention are selected from the group consisting of:

15 (a) the DNA sequence of Figure 1 (SEQ ID NO:1);
 (b) the DNA sequence of Figure 8 (SEQ ID NO:3);
 (c) the DNA sequence of Figure 9 (SEQ ID NO:5);
 (d) the DNA sequence of Figure 10 (SEQ ID NO:7);
 (e) analogues or derivatives of the foregoing DNA
20 sequences;

(f) DNA sequences degenerate to any of the foregoing DNA sequences; and

(g) fragments of any of the foregoing DNA sequences; wherein said sequences encode a product that displays the
25 immunological activity of the *Neisseria meningitidis* 22 kDa surface protein.

Such fragments are preferably peptides as depicted in Figure 15 (SEQ ID NO:9, through SEQ ID NO:26).

Preferably, this invention also provides, for the
30 first time, a DNA sequence coding for the *Neisseria meningitidis* 22 kDa surface protein (SEQ ID NO:1). More preferred DNA sequences of this invention are selected from the group consisting of:

(a) the DNA sequence of Figure 1 (SEQ ID NO:1);

(b) analogues or derivatives of the foregoing DNA sequences;

(c) DNA sequences degenerate to any of the foregoing DNA sequences; and

5 (d) fragments of any of the foregoing DNA sequences; wherein said sequences encode a product that displays the immunological activity of the *Neisseria meningitidis* 22 kDa surface protein.

Analogues and derivatives of the *Neisseria*
10 *meningitidis* 22 kDa surface protein coding gene will hybridize to the 22 kDa surface protein-coding gene under the conditions described in Example 4.

For purposes of this invention, the fragments, analogues and derivatives of the *Neisseria meningitidis* 22
15 kDa surface protein have the "immunological activity" of the *Neisseria meningitidis* 22 kDa surface protein if they can induce, when injected into an animal host, the development of an immunological response that can protect against *Neisseria meningitidis* infection. One of skill in
20 the art may determine whether a particular DNA sequence encodes a product that displays the immunological activity of the *Neisseria meningitidis* 22 kDa surface protein by following the procedures set forth herein in Example 6.

The *Neisseria meningitidis* surface proteins of this
25 invention may be isolated by a method comprising the following steps:

a) isolating a culture of *Neisseria meningitidis* bacteria,

b) isolating an outer membrane portion from the
30 culture of the bacteria; and

c) isolating said antigen from the outer membrane portion.

In particular, the foregoing step (c) may include the additional steps of treating the *Neisseria meningitidis*
35 outer membrane protein extracts with proteinase K,

followed by protein fractionation using conventional separation techniques such as ion exchange and gel chromatography and electrophoresis.

Alternatively and preferably, the *Neisseria meningitidis* surface proteins of this invention may be produced by the use of molecular biology techniques, as more particularly described in Example 3 herein. The use of molecular biology techniques is particularly well-suited for the preparation of substantially pure recombinant *Neisseria meningitidis* 22 kDa surface protein.

Thus according to a further aspect of the invention we provide a process for the production of recombinant *Neisseria meningitidis* 22 kDa surface protein, including fragments, analogues and derivatives thereof, comprising the steps of (1) culturing a unicellular host organism transformed with a recombinant DNA molecule including a DNA sequence coding for said protein, fragment, analogue or derivative and one or more expression control sequences operatively linked to the DNA sequence, and (2) recovering a substantially pure protein, fragment, analogue or derivative.

As is well known in the art, in order to obtain high expression levels of a transfected gene in a host, the gene must be operatively linked to transcriptional and translational expression control sequences that are functional in the chosen expression host. Preferably, the expression control sequences, and the gene of interest, will be contained in an expression vector that further comprises a bacterial selection marker and origin of replication. If the expression host is a eukaryotic cell, the expression vector should further comprise an expression marker useful in the expression host.

A wide variety of expression host/vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression vectors for eukaryotic hosts

include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Useful expression vectors for bacterial hosts include known bacterial plasmids, such as plasmids from *E.coli*, including col E1, pCR1, pBR322, pMB9 and their derivatives, wider host range plasmids, such as RP4, phage DNAs, e.g., the numerous derivatives of phage lambda, e.g. NM989, and other DNA phages, such as M13 and filamentous single stranded DNA phages. Useful expression vectors for yeast cells include the 2 mu plasmid and derivatives thereof. Useful vectors for insect cells include pVL 941.

In addition, any of a wide variety of expression control sequences may be used in these vectors to express the DNA sequences of this invention. Such useful expression control sequences include the expression control sequences associated with structural genes of the foregoing expression vectors. Examples of useful expression control sequences include, for example, the early and late promoters of SV40 or adenovirus, the lac system, the trp system, the TAC or TRC system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast alpha-mating system and other sequences known to control expression of genes of prokaryotic and eukaryotic cells or their viruses, and various combinations thereof. The *Neisseria meningitidis* 22 kDa surface protein's expression control sequence is particularly useful in the expression of the *Neisseria meningitidis* 22 kDa surface protein in *E.coli* (Example 3).

Host cells transformed with the foregoing vectors form a further aspect of this invention. A wide variety of unicellular host cells are useful in expressing the DNA

sequences of this invention. These hosts may include well known eukaryotic and prokaryotic hosts, such as strains of *E.coli*, *Pseudomonas*, *Bacillus*, *Streptomyces*, fungi, yeast, insect cells such as *Spodoptera frugiperda* (SF9), animal
5 cells such as CHO and mouse cells, African green monkey cells such as COS 1, COS 7, BSC 1, BSC 40, and BMT 10, and human cells and plant cells in tissue culture. Preferred host organisms include bacteria such as *E.coli* and *Bacillus subtilis* and mammalian cells in tissue culture.

10 It should of course be understood that not all vectors and expression control sequences will function equally well to express the DNA sequences of this invention. Neither will all hosts function equally well with the same expression system. However, one of skill in
15 the art may make a selection among these vectors, expression control sequences and hosts without undue experimentation and without departing from the scope of this invention. For example, in selecting a vector, the host must be considered because the vector must replicate
20 in it. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered.

In selecting an expression control sequence, a
25 variety of factors should also be considered. These include, for example, the relative strength of the sequence, its controllability, and its compatibility with the DNA sequences of this invention, particularly as regards potential secondary structures. Unicellular hosts
30 should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for by the DNA sequences of this invention, their secretion characteristics, their ability to fold the protein correctly, their fermentation or culture
35 requirements, and the ease of purification from them of

the products coded for by the DNA sequences of this invention.

Within these parameters, one of skill in the art may select various vector/expression control sequence/host
5 combinations that will express the DNA sequences of this invention on fermentation or in large scale animal culture.

The polypeptides encoded by the DNA sequences of this invention may be isolated from the fermentation or cell
10 culture and purified using any of a variety of conventional methods. One of skill in the art may select the most appropriate isolation and purification techniques without departing from the scope of this invention.

The *Neisseria meningitidis* surface proteins of this
15 invention are useful in prophylactic, therapeutic and diagnostic compositions for preventing, treating and diagnosing diseases caused by *Neisseria meningitidis* infection.

The *Neisseria meningitidis* surface proteins of this
20 invention are useful in prophylactic, therapeutic and diagnostic compositions for preventing, treating and diagnosing diseases caused by *Neisseria gonorrhoeae*, or *Neisseria lactamica* infection.

The *Neisseria meningitidis* surface proteins according
25 to this invention are particularly well-suited for the generation of antibodies and for the development of a protective response against *Neisseria meningitidis* diseases.

The *Neisseria meningitidis* surface proteins according
30 to this invention are particularly well-suited for the generation of antibodies and for the development of a protective response against *Neisseria gonorrhoeae* or *Neisseria lactamica* diseases.

In particular, we provide a *Neisseria meningitidis* 22
35 kDa surface protein having an amino acid sequence of

Figure 1 (SEQ ID NO:2) or a fragment, analogue, or derivative thereof for use as an immunogen and as a vaccine.

5 In particular, we provide a *Neisseria meningitidis* 22 kDa surface protein having an amino acid sequence of Figure 1 (SEQ ID NO:2), Figure 8 (SEQ ID NO:4), Figure 9 (SEQ ID NO:6), or Figure 10 (SEQ ID NO:8), or a fragment, analogue, or derivative thereof for use as an immunogen and as a vaccine.

10 Standard immunological techniques may be employed with the *Neisseria meningitidis* surface proteins in order to use them as immunogens and as vaccines. In particular, any suitable host may be injected with a pharmaceutically effective amount of the *Neisseria meningitidis* 22 kDa
15 surface protein to generate monoclonal or polyvalent anti-*Neisseria meningitidis* antibodies or to induce the development of a protective immunological response against *Neisseria meningitidis* diseases. Prior to injection of the host, the *Neisseria meningitidis* surface proteins may
20 be formulated in a suitable vehicle, and thus we provide a pharmaceutical composition comprising one or more *Neisseria meningitidis* surface antigens or fragments thereof. Preferably, the antigen is the *Neisseria meningitidis* 22 kDa surface protein or fragments,
25 analogues or derivatives thereof together with one or more pharmaceutically acceptable excipients. As used herein, "pharmaceutically effective amount" refers to an amount of one or more *Neisseria meningitidis* surface antigens or fragments thereof that elicits a sufficient titer of anti-
30 *Neisseria meningitidis* antibodies to treat or prevent *Neisseria meningitidis* infection.

The *Neisseria meningitidis* surface proteins of this invention may also form the basis of a diagnostic test for *Neisseria meningitidis* infection. Several diagnostic
35 methods are possible. For example, this invention

provides a method for the detection of *Neisseria meningitidis* antigen in a biological sample containing or suspected of containing *Neisseria meningitidis* antigen comprising:

- 5 a) isolating the biological sample from a patient;
- b) incubating an anti-*Neisseria meningitidis* 22 kDa surface protein antibody or fragment thereof with the biological sample to form a mixture; and
- c) detecting specifically bound antibody or bound
10 fragment in the mixture which indicates the presence of *Neisseria meningitidis* antigen.

Preferred antibodies in the foregoing diagnostic method are Me-1 and Me-7.

Alternatively, this invention provides a method for
15 the detection of antibody specific to *Neisseria meningitidis* antigen in a biological sample containing or suspected of containing said antibody comprising:

- a) isolating the biological sample from a patient;
 - b) incubating a *Neisseria meningitidis* surface
20 protein of this invention or fragment thereof with the biological sample to form a mixture; and
 - c) detecting specifically bound antigen or bound fragment in the mixture which indicates the presence of antibody specific to *Neisseria meningitidis* antigen.
- 25 One of skill in the art will recognize that this diagnostic test may take several forms, including an immunological test such as an enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay or a latex agglutination assay, essentially to determine whether antibodies
30 specific for the protein are present in an organism.

The DNA sequences of this invention may also be used to design DNA probes for use in detecting the presence of the pathogenic *Neisseria* bacteria in a biological suspected of containing such bacteria. The detection
35 method of this invention comprises the steps of:

a) isolating the biological sample from a patient;
b) incubating a DNA probe having a DNA sequence of this invention with the biological sample to form a mixture; and

5 c) detecting specifically bound DNA probe in the mixture which indicates the presence of *Neisseria* bacteria.

Preferred DNA probes have the base pair sequence of Figure 1 (SEQ ID NO:1), Figure 8 (SEQ ID NO:3), Figure 9
10 (SEQ ID NO:5), or Figure 10 (SEQ ID NO:7), or consensus sequence of Figure 11 (SEQ ID NO:9).

A more preferred DNA probe has the 525 base pair sequence of Figure 1 (SEQ ID NO:1).

The DNA probes of this invention may also be used for
15 detecting circulating *Neisseria meningitidis* nucleic acids in a sample, for example using a polymerase chain reaction, as a method of diagnosing *Neisseria meningitidis* infections. The probe may be synthesized using conventional techniques and may be immobilized on a solid
20 phase, or may be labeled with a detectable label.

A preferred DNA probe for this application is an oligomer having a sequence complementary to at least about 6 contiguous nucleotides of the *Neisseria meningitidis* 22
kDa surface protein gene of Figure 1 (SEQ ID NO:1), Figure
25 8 (SEQ ID NO:3), Figure 9 (SEQ ID NO:5), Figure 10 (SEQ ID NO:7), or consensus sequence of Figure 11 (SEQ ID NO:9).

A more preferred DNA probe for this application is an oligomer having a sequence complementary to at least about 6 contiguous nucleotides of the *Neisseria meningitidis* 22
30 kDa surface protein gene of Figure 1 (SEQ ID NO:1).

Another diagnostic method for the detection of *Neisseria meningitidis* in a patient comprises the steps of:

a) labeling an antibody of this invention or
35 fragment thereof with a detectable label;

b) administering the labeled antibody or labeled fragment to the patient; and

c) detecting specifically bound labeled antibody or labeled fragment in the patient which indicates the presence of *Neisseria meningitidis*.

For purification of any anti-*Neisseria meningitidis* surface protein antibody, use may be made of affinity chromatography employing an immobilized *Neisseria meningitidis* surface protein as the affinity medium.

Thus according to another aspect of the invention we provide a *Neisseria meningitidis* 22 kDa surface protein having an amino acid sequence which includes the sequence of Figure 1 (SEQ ID NO:2), Figure 8 (SEQ ID NO:4), Figure 9 (SEQ ID NO:6), or Figure 10 (SEQ ID NO:8), or portion thereof or an analogue thereof, covalently bound to an insoluble support.

Thus according to a preferred aspect of the invention we provide a *Neisseria meningitidis* 22 kDa surface protein having an amino acid sequence which includes the sequence of Figure 1 (SEQ ID NO:2), or portion thereof or an analogue thereof, covalently bound to an insoluble support.

A further feature of the invention is the use of the *Neisseria meningitidis* surface proteins of this invention as immunogens for the production of specific antibodies for the diagnosis and in particular the treatment of *Neisseria meningitidis* infection. Suitable antibodies may be determined using appropriate screening methods, for example by measuring the ability of a particular antibody to passively protect against *Neisseria meningitidis* infection in a test model. One example of an animal model is the mouse model described in the Examples herein. The antibody may be a whole antibody or an antigen-binding fragment thereof and may in general belong to any immunoglobulin class. The antibody or fragment may be of

animal origin, specifically of mammalian origin and more specifically of murine, rat or human origin. It may be a natural antibody or a fragment thereof, or if desired, a recombinant antibody or antibody fragment. The term
5 recombinant antibody or antibody fragment means antibody or antibody fragment which were produced using molecular biology techniques. The antibody or antibody fragments may be of polyclonal, or preferentially, monoclonal origin. It may be specific for a number of epitopes
10 associated with the *Neisseria meningitidis* surface proteins but it is preferably specific for one. Preferably, the antibody or fragments thereof will be specific for one or more epitopes associated with the *Neisseria meningitidis* 22 kDa surface protein. Also
15 preferred are the monoclonal antibodies Me-1 and Me-7 described herein.

EXAMPLES

In order that this invention may be better
20 understood, the following examples are set forth. These examples are for purposes of illustration only, and are not to be construed as limiting the scope of the invention in any manner.

Example 1 describes the treatment of *Neisseria*
25 *meningitidis* outer membrane preparation with proteolytic enzymes and the subsequent identification of the *Neisseria meningitidis* 22 kDa surface protein.

Example 2 describes the preparation of monoclonal antibodies specific for *Neisseria meningitidis* 22 kDa
30 surface protein.

Example 3 describes the preparation of *Neisseria meningitidis* recombinant 22 kDa surface protein.

Example 4 describes the use of DNA probes for the identification of organisms expressing the *Neisseria*
35 *meningitidis* 22 kDa surface protein.

Example 5 describes the use of an anti-*Neisseria meningitidis* 22 kDa surface protein monoclonal antibody to protect mice against *Neisseria meningitidis* infection.

Example 6 describes the use of purified recombinant 22 kDa surface protein to induce a protective response against *Neisseria meningitidis* infection.

Example 7 describes the identification of the sequence for the 22kDa protein and protein-coding gene for other strains of *Neisseria meningitidis* (MCH88, and Z4063), and one strain of *Neisseria gonorrhoeae*.

Example 8 describes the immunological response of rabbits and monkeys to the 22kDa *Neisseria meningitidis* surface protein.

Example 9 describes the procedure used to map the different immunological epitopes of the 22kDa *Neisseria meningitidis* surface protein.

Example 10 describes the induction by heat of an expression vector for the large scale production of the 22 kDa surface protein.

Example 11 describes a purification process for the 22kDa surface protein when produced by recombinant technology.

Example 12 describes the use of 22kDa surface protein as a human vaccine.

EXAMPLE 1 Treatment Of *Neisseria meningitidis* Outer Membrane Preparations With Proteolytic Enzymes And The Subsequent Identification Of An Enzyme Resistant *Neisseria meningitidis* 22 kDa Surface Protein

Several antigenic preparations derived from whole cell, lithium chloride, or sarcosyl extracts were used to study the ultrastructure of *Neisseria meningitidis* outer membrane. The outer membrane of Gram-negative bacteria

acts as an interface between the environment and the interior of the cell and contains most of the antigens that are freely exposed to the host immune defense. The main goal of the study was the identification of new
5 antigens which can induce a protective response against *Neisseria meningitidis*. One approach used by the inventors to identify such antigens, was the partial disruption of the antigenic preparations mentioned above with proteolytic enzymes. The antigenic determinants
10 generated by the enzymatic treatments could then be identified by the subsequent analysis of the immunological and protective properties of these treated antigenic preparations. To our surprise we observed after electrophoretic resolution of *Neisseria meningitidis*
15 lithium chloride outer membrane extracts, that one low molecular weight band, which was stained with Coomassie Brilliant Blue R-250, was not destroyed by proteolytic enzyme treatments. Coomassie Blue is used to stain proteins and peptides and has no or very little affinity
20 for the polysaccharides or lipids which are also key components of the outer membrane. The fact that this low molecular weight antigen was stained by Coomassie blue suggested that at least part of it is made up of polypeptides that are not digested by proteolytic enzymes,
25 or that are protected against the action of the enzymes by other surface structures. Moreover, as demonstrated below the very potent enzyme proteinase K did not digest this low molecular weight antigen even after extensive treatments.

30 Lithium chloride extraction was used to obtain the outer membrane preparations from different strains of *Neisseria meningitidis* and was performed in a manner previously described by the inventors [Brodeur et al., Infect. Immun., 50, p. 510 (1985)]. The protein content
35 of these preparations were determined by the Lowry method

adapted to membrane fractions [Lowry et al., J. Biol. Chem. 193, p. 265 (1951)]. Outer membrane preparations derived from *Neisseria meningitidis* strain 608B (B:2a:P1.2) were treated for 24 hours at 37°C and

5 continuous shaking with either α -chymotrypsin from bovine pancreas (E.C. 3.4.21.1) (Sigma) or trypsin type 1 from bovine pancreas (E.C. 3.4.21.4) (Sigma). The enzyme concentration was adjusted at 2 mg per mg of protein to be treated. The same outer membrane preparations were also

10 treated with different concentrations (0.5 to 24 mg per mg of protein) of Proteinase K from *Tritirachium album limber* (Sigma or Boehringer Mannheim, Laval, Canada) (E.C. 3.4.21.14). In order to promote protein digestion by proteinase K, different experimental conditions were used.

15 The samples were incubated for 1 hour, 2 hours, 24 hours or 48 hours at 37°C or 56°C with or without shaking. The pH of the mixture samples was adjusted at either pH 7.2 or pH 9.0. One % (vol/vol) of sodium dodecyl sulfate (SDS) was also added to certain samples. Immediately after

20 treatment the samples were resolved by SDS-PAGE gel electrophoresis using the MiniProteanII® (Bio-Rad, Mississauga, Ontario, Canada) system on 14% (wt/vol) gels according to the manufacturer's instructions. Proteins were heated to 100°C for 5 minutes with 2-mercaptoethanol

25 and SDS, separated on 14% SDS gels, and stained with Coomassie Brilliant Blue R-250.

Figure 2 presents the migration profile on 14% SDS-PAGE gel of the proteins present in outer membrane preparations derived from *Neisseria meningitidis* strain

30 608B (B:2a:P1.2) after treatment at 37°C for 24 hours with α -chymotrypsin and trypsin. Extensive proteolytic digestion of the high molecular weight proteins and of several major outer membrane proteins can be observed for the treated samples (Figure 2, lanes 3 and 4) compared to

35 the untreated control (Figure 2, lane 2). On the

contrary, a protein band with an apparent molecular weight of 22 kDa was not affected even after 24 hours of contact with either proteolytic enzyme.

This unique protein was further studied using a more aggressive proteolytic treatment with Proteinase K (Figure 3). Proteinase K is one of the most powerful proteolytic enzymes since it has a low peptide bond specificity and wide pH optimum. Surprisingly, the 22 kDa protein was resistant to digestion by 2 International Units (IU) of proteinase K for 24 hours at 56°C (Figure 3, lane 2). This treatment is often used in our laboratory to produce lipopolysaccharides or DNA that are almost free of proteins. Indeed, only small polypeptides can be seen after such an aggressive proteolytic treatment of the outer membrane preparation. Furthermore, longer treatments, up to 48 hours, or higher enzyme concentrations (up to 24 IU) did not alter the amount of the 22 kDa protein. The amount and migration on SDS-PAGE gel of the 22 kDa protein were not affected when the pH of the reaction mixtures was increased to pH 9.0, or when 1.0% of SDS, a strong protein denaturant was added (Figure 3, lanes 4, 6 and 8). The combined use of these two denaturing conditions would normally result in the complete digestion of the proteins present in the outer membrane preparations, leaving only amino acid residues. Polypeptides of low molecular weight were often observed in the digests and were assumed to be fragments of sensitive proteins not effectively digested during the enzymatic treatments. These fragments were most probably protected from further degradation by the carbohydrates and lipids present in the outer membrane. The bands with apparent molecular weight of 28 kDa and 34 kDa which are present in treated samples are respectively the residual enzyme and a contaminating protein present in all enzyme preparations tested.

Interestingly, this study about the resistance of the 22kDa protein to proteases indicated that another protein band with apparent molecular weight of 18kDa seems to be also resistant to enzymatic degradation (Figure 3a). Clues about this 18kDa protein band were obtained when the migration profiles on SDS-PAGE gels of affinity purified recombinant 22kDa protein were analyzed (Figure 3b). The 18 kDa band was apparent only when the affinity purified recombinant 22kDa protein was heated for an extended period of time in sample buffer containing the detergent SDS before it was applied on the gel. N-terminal amino acid analysis using the Edman degradation (Example 3) clearly established that the amino acid residues (E-G-A-S-G-F-Y-V-Q) identified on the 18 kDa band corresponded to the amino acids 1-9 (SEQ ID NO:1). These results indicate that the 18 and 22kDa bands as seen on the SDS-PAGE is in fact derived from the same protein. This last result also indicates that the leader sequence is cleaved from the mature 18 kDa protein. Further studies will be done to identify the molecular modifications explaining this shift in apparent molecular weight and to evaluate their impact on the antigenic and protective properties of the protein.

In conclusion, the discovery of a *Neisseria meningitidis* outer membrane protein with the very rare property of being resistant to proteolytic digestion warranted further study of its molecular and immunological characteristics. The purified recombinant 22 kDa surface protein produced by *Escherichia coli* in Example 3 is also highly resistant to proteinase K digestion. We are presently trying to understand the mechanism which confers to the *Neisseria meningitidis* 22 kDa surface protein this unusual resistance to proteolytic enzymes.

**EXAMPLE 2 Generation of Monoclonal Antibodies Specific for
the 22 kDa *Neisseria meningitidis* Surface
Protein**

5 The monoclonal antibodies described herein were
obtained from three independent fusion experiments.
Female Balb/c mice (Charles River Laboratories, St-
Constant, Québec, Canada) were immunized with outer
membrane preparations obtained from *Neisseria meningitidis*
10 strains 604A, 608B and 2241C respectively serogrouped A, B
and C. The lithium chloride extraction used to obtain
these outer membrane preparations was performed in a
manner previously described by the inventors. [Brodeur
et al., Infect. Immun. 50, p. 510 (1985)]. The protein
15 content of these preparations were determined by the Lowry
method adapted to membrane fractions [Lowry et al., J.
Biol. Chem. 193, p. 265 (1951)]. Groups of mice were
injected intraperitoneally or subcutaneously twice, at
three-week intervals with 10 mg of different combinations
20 of the outer membrane preparations described above.
Depending on the group of mice, the adjuvants used for the
immunizations were either Freund's complete or incomplete
adjuvant (Gibco Laboratories, Grand Island, N.Y.), or
QuilA (CedarLane Laboratories, Hornby, Ont., Canada).
25 Three days before the fusion procedure, the immunized mice
received a final intravenous injection of 10 mg of one of
the outer membrane preparations described above. The
fusion protocol used to produce the hybridoma cell lines
secreting the desired monoclonal antibody was described
30 previously by the inventors [Hamel et al., J. Med.
Microbiol., 25, p. 2434 (1987)]. The class, subclass and
light-chain type of monoclonal antibodies Me-1, Me-2, Me-
3, Me-5, Me-6 and Me-7 were determined by ELISA as
previously reported [Martin et al., J. Clin. Microbiol.,
35 28, p. 1720 (1990)] and are presented in Table 1.

The specificity of the monoclonal antibodies was established using Western immunoblotting following the method previously described by the inventors [Martin et al., Eur. J. Immunol. 18, p. 601 (1988)] with the following modifications. Outer membrane preparations obtained from different strains of *Neisseria meningitidis* were resolved on 14% SDS-PAGE gels. The proteins were transferred from the gels to nitrocellulose membranes using a semi-dry apparatus (Bio-Rad). A current of 60 mA per gel (6X10cm) was applied for 10 minutes in the electroblot buffer consisting of 25 mM Tris-HCl, 192 mM glycine and 20% (vol/vol) methanol, pH 8.3. The Western immunoblotting experiments clearly indicated that the monoclonal antibodies Me-1, Me-2, Me-3, Me-5, Me-6 and Me-7 recognized their specific epitopes on the *Neisseria meningitidis* 22 kDa protein (Figure 4A). Analysis of the SDS-PAGE gels and the corresponding Western immunoblots also indicated that the apparent molecular weight of this protein does not vary from one strain to another. However, the amount of protein present in the outer membrane preparations varied from one strain to another and was not related to the serogroup of the strain. Moreover, these monoclonal antibodies still recognized their epitopes on the *Neisseria meningitidis* 22 kDa surface protein after treatment of the outer membrane preparation with 2 IU of proteinase K per mg of protein (treatment described in Example 1, supra) (Figure 4B). Interestingly, the epitopes remained intact after the enzyme digestion thus confirming that even if they are accessible in the membrane preparation to the antibodies they are not destroyed by the enzyme treatment. This latter result suggested that the mechanism which explains the observed proteinase K resistance is most probably not related to surface structures blocking the access of the enzyme to the protein, or to the protection offered by the

membrane to proteins which are deeply embedded. While not shown in Figure 4, the results of the immunoblots for Me-1 were consistent with the results for the other five monoclonal antibodies.

- 5 A series of experiments were performed to partially characterize the *Neisseria meningitidis* 22 kDa surface protein and to differentiate it from the other known meningococcal surface proteins. No shift in apparent molecular weight on SDS-PAGE gel of the *Neisseria*
- 10 *meningitidis* 22 kDa surface protein was noted when outer membrane preparations were heated at 100°C for 5 minutes, or at 37°C and 56°C for 30 minutes in electrophoresis sample buffer with or without 2-mercaptoethanol. This indicated that the migration of the 22kDa surface protein,
- 15 when present in the outer membrane, was not heat- or 2-mercaptoethanol-modifiable.

- Sodium periodate oxidation was used to determine if the monoclonal antibodies reacted with carbohydrate epitopes present in the outer membrane preparations
- 20 extracted from *Neisseria meningitidis* organisms. The method used to perform this experiment was previously described by the inventors. [Martin et al., Infect. Immun., 60, pp. 2718-2725 (1992)]. Treatment of outer membrane preparations with 100 mM of sodium periodate for
- 25 1 hour at room temperature did not alter the reactivity of the monoclonal antibodies toward the *Neisseria meningitidis* 22 kDa surface protein. This treatment normally abolishes the binding of antibodies that are specific for carbohydrates.

- 30 Monoclonal antibody 2-1-CA2 (provided by Dr. A. Bhattacharjee, Walter Reed Army Institute of Research, Washington, D.C.) is specific for the lip protein (also called H.8), a surface antigen common to all pathogenic *Neisseria* species. The reactivity of this monoclonal
- 35 antibody with outer membrane preparations was compared to

the reactivity of monoclonal antibody Me-5. The lip-specific monoclonal antibody reacted with a protein band having an apparent molecular weight of 30 kDa, while monoclonal antibody Me-5 reacted with the protein band of 22 kDa. This result clearly indicates that there is no relationship between *Neisseria meningitidis* 22 kDa surface protein and the lip protein, another highly conserved outer membrane protein.

To verify the exposure of the 22 kDa protein at the surface of intact *Neisseria meningitidis* bacterial cells, a radioimmunoassay was performed as previously described by the inventors [Proulx et al., Infec. Immun., 59, p. 963 (1991)]. Six-hour and 18-hour bacterial cultures were used for this assay. The six monoclonal antibodies were reacted with 9 *Neisseria meningitidis* strains (the serogroup of the strain is indicated in parentheses on Figure 5), 2 *Neisseria gonorrhoeae* strains ("NG"), 2 *Moraxella catarrhalis* strains ("MC") and 2 *Neisseria lactamica* strains ("NL"). The radioimmunoassay confirmed that the epitopes recognized by the monoclonal antibodies are exposed at the surface of intact *Neisseria meningitidis* isolates of different serotypes and serogroups and should also be accessible to the proteolytic enzymes (Figure 5). The monoclonal antibodies bound strongly to their target epitopes on the surface of all *Neisseria meningitidis* strains tested. The recorded binding values (between 3,000 to 35,000 CPM), varied from one strain to another, and with the physiological state of the bacteria. A *Haemophilus influenzae* porin-specific monoclonal antibody was used as a negative control for each bacterial strain. Counts below 500 CPM were obtained and subsequently subtracted from each binding value. With respect to the *Neisseria meningitidis* strains tested in this assay, the results shown in Figure 5 for monoclonal antibodies Me-5 and Me-7 are representative of the results

obtained with monoclonal antibodies Me-1, Me-2, Me-3 and Me-6. With respect to the other bacterial strains tested, the binding activities shown for Me-7 are representative of the binding activities obtained with other monoclonal
5 antibodies that recognized the same bacterial strain.

The antigenic conservation of the epitopes recognized by the monoclonal antibodies was also evaluated. A dot enzyme immunoassay was used for the rapid screening of the monoclonal antibodies against a large number of bacterial
10 strains. This assay was performed as previously described by the inventors [Lussier et al., *J. Immunoassay*, 10, p. 373 (1989)]. A collection of 71 *Neisseria meningitidis* strains was used in this study. The sample included 19 isolates of serogroup A, 23 isolates of serogroup B, 13
15 isolates of serogroup C, 1 isolate of serogroup 29E, 6 isolates of serogroup W-135, 1 isolate of serogroup X, 2 isolates of serogroup Y, 2 isolates of serogroup Z, and 4 isolates that were not serogrouped ("NS"). These isolates were obtained from the Caribbean Epidemiology Centre, Port
20 of Spain, Trinidad; Children's Hospital of Eastern Ontario, Ottawa, Canada; Department of Saskatchewan Health, Regina, Canada; Laboratoire de Santé Publique du Québec, Montreal, Canada; Max-Planck Institut für Molekulare Genetik, Berlin, FRG; Montreal Children
25 Hospital, Montreal, Canada; Victoria General Hospital, Halifax, Canada; and our own strains collection. The following bacterial species were also tested: 16 *Neisseria gonorrhoeae*, 4 *Neisseria cinerea*, 5 *Neisseria lactamica*, 1 *Neisseria flava*, 1 *Neisseria flavescens*, 3 *Neisseria*
30 *mucosa*, 4 *Neisseria perflava/sicca*, 4 *Neisseria perflava*, 1 *Neisseria sicca*, 1 *Neisseria subflava* and 5 *Moraxella catarrhalis*, 1 *Alcaligenes faecalis* (ATCC 8750), 1 *Citrobacter freundii* (ATCC 2080), 1 *Edwardsiella tarda* (ATCC 15947), 1 *Enterobacter cloaca* (ATCC 23355), 1
35 *Enterobacter aerogenes* (ATCC 13048), 1 *Escherichia coli*, 1

Flavobacterium odoratum, 1 *Haemophilus influenzae* type b (Eagan strain), 1 *Klebsiella pneumoniae* (ATCC 13883), 1 *Proteus rettgeri* (ATCC 25932), 1 *Proteus vulgaris* (ATCC 13315), 1 *Pseudomonas aeruginosa* (ATCC 9027), 1 *Salmonella* 5 *typhimurium* (ATCC 14028), 1 *Serratia marcescens* (ATCC 8100), 1 *Shigella flexneri* (ATCC 12022), 1 *Shigella sonnei* (ATCC 9290). They were obtained from the American Type Culture Collection or a collection held in the Laboratory Centre for Disease Control, Ottawa, Canada. The 10 reactivities of the monoclonal antibodies with the most relevant *Neisseria* strains are presented in Table 1. One monoclonal antibody, Me-7, recognized its specific epitope on 100% of the 71 *Neisseria meningitidis* strains tested. This monoclonal antibody, as well as Me-2, Me-3, Me-5 and 15 Me-6 also reacted with certain strains belonging to other *Neisserial* species indicating that their specific epitope is also expressed by other closely related *Neisseriaceae*.

Except for a faint reaction with one *Neisseria lactamica* strain, monoclonal antibody Me-1 reacted only with 20 *Neisseria meningitidis* isolates. Me-1 was further tested with another sample of 177 *Neisseria meningitidis* isolates and was able to correctly identify more than 99% of the total *Neisseria meningitidis* strains tested. Besides the *Neisseria* strains presented in Table 1, the 25 monoclonal antibodies did not react with any of the other bacterial species mentioned above.

In conclusion, six monoclonal antibodies which specifically reacted with the *Neisseria meningitidis* 22 kDa surface protein were generated by the inventors. 30 Using these monoclonal antibodies we demonstrated that their specific epitopes are 1) located on a proteinase K resistant 22 kDa protein present in the outer membrane of *Neisseria meningitidis*, 2) conserved among *Neisseria meningitidis* isolates, 3) exposed at the surface of intact 35 *Neisseria meningitidis* cells and accessible to antibody,

and 4) the reactivity of these monoclonal antibodies with the *Neisseria meningitidis* 22 kDa surface protein is not modified by a treatment with sodium periodate, suggesting that their specific epitopes are not located on
5 carbohydrates.

Although we found that the migration of the *Neisseria meningitidis* 22kDa protein is moved to an apparent molecular weight of about 18kDa when heated under stringent conditions, we observed that the migration is
10 not modified by 2-mercaptoethanol treatment.

We also demonstrated that the *Neisseria meningitidis* 22 kDa surface protein has no antigenic similarity with the lip protein, another low molecular weight and highly conserved protein present in the outer membrane of
15 *Neisseria meningitidis*.

As will be presented in Example 3, these monoclonal antibodies also reacted with the purified, recombinant 22 kDa surface protein produced after transformation of *Escherichia coli* strain BL21 (DE3) with a plasmid vector
20 pNP2202 containing the gene coding for the *Neisseria meningitidis* 22 kDa surface protein.

Table 1. R activity of the monoclonal antibodies with *Neisseria* isolates

Number of <i>Neisseria</i> isolates recognized by the monoclonal antibodies														
Name	Isotype	Serogroup of <i>Neisseria meningitidis</i>									Total (71)	<i>Neisseria</i> <i>catarrhalis</i> (5)	<i>Neisseria</i> <i>gonorrhoeae</i> (16)	<i>Neisseria</i> <i>lactamica</i> (5)
		A (19)	B (23)	C (13)	29B (1)	W135 (6)	X (1)	Y (2)	Z (2)	MS ¹ (4)				
Me-1	IgG2a (K)	19	22	13	1	6	1	2	2	3	69	0	0	1
Me-2	IgG2a (K)	19	20	13	1	6	0	2	2	4	67	0	2	0
Me-3	IgG3 (K)	19	22	13	1	6	1	2	2	3	69	0	2	4
Me-5	IgG2a (K)	19	22	13	1	6	1	2	2	3	69	0	2	0
Me-6	IgG1 (K)	19	23	13	1	6	1	2	2	3	70	0	2	4
Me-7	IgG2a (K)	19	23	13	1	6	1	2	2	4	71	5	2	4

¹isolates not serogrouped

5

EXAMPLE 3 Molecular Cloning, Sequencing Of The Gene, High Yield Expression And Purification Of The *Neisseria meningitidis* 22 kDa Surface Protein

10 **A. Molecular Cloning**

A LambdaGEM-11 genomic DNA library from *Neisseria meningitidis* strain 608B (B:2a:P1.2) was constructed according to the manufacturer's recommendations (Promega CO, Madison, WI). Briefly, the genomic DNA of the 608B strain was partially digested with Sau 3AI, and fragments ranging between 9 and 23 Kb were purified on agarose gel before being ligated to the Bam HI sites of the LambdaGEM-11 arms. The resulting recombinant phages were used to infect *Escherichia coli* strain LE392 (Promega) which was then plated onto LB agar plates. Nineteen positive plaques were identified after the immuno-screening of the library with the *Neisseria meningitidis* 22 kDa surface protein-specific monoclonal antibodies of Example 2 using the following protocol. The plates were incubated 15

minutes at -20°C to harden the top agar. Nitrocellulose filters were gently applied onto the surface of the plates for 30 minutes at 4°C to absorb the proteins produced by the recombinant viral clones. The filters were then

5 washed in PBS-Tween 0.02% (vol/vol) and immunoblotted as described previously [Lussier et al., J. Immunoassay, 10, p. 373 (1989)]. After amplification and DNA purification, one viral clone, designated clone 8, which had a 13 Kb insert was selected for the subcloning experiments. After

10 digestion of this clone with Sac I, two fragments of 5 and 8 Kb were obtained. These fragments were purified on agarose gel and ligated into the Sac I restriction site of the low copy number plasmid pWKS30 [Wang and Kushner, Gene, 100, p. 195 (1991)]. The recombinant plasmids were

15 used to transform *Escherichia coli* strain JM109 (Promega) by electroporation (Bio-Rad, Mississauga, Ont., Canada) following the manufacturer's recommendations, and the resulting colonies were screened with the *Neisseria meningitidis* 22 kDa surface protein-specific monoclonal

20 antibodies of Example 2. Positive colonies were observed only when the bacteria were transformed with the plasmid carrying the 8 Kb insert. Western blot analysis (the methodology was described in Example 2) of the positive clones showed that the protein expressed by *Escherichia*

25 *coli* was complete and migrated on SDS-PAGE gel like the *Neisseria meningitidis* 22 kDa surface protein. To further reduce the size of the insert, a clone containing the 8 Kb fragment was digested with Cla I and a 2.75 Kb fragment was then ligated into the Cla I site of the pWKS30

30 plasmid. Western blot analysis of the resulting clones clearly indicated once again that the protein expressed by *Escherichia coli* was complete and migrated on SDS-PAGE gel like the native *Neisseria meningitidis* 22 kDa surface protein.

After restriction analysis, two clones, designated pNP2202 and pNP2203, were shown to carry the 2.75 Kb insert in opposite orientations and were selected to proceed with the sequencing of the gene coding for the

5 *Neisseria meningitidis* 22 kDa surface protein. The "Double Stranded Nested Deletion Kit" from Pharmacia Biotech Inc. (Piscataway, NJ) was used according to the manufacturer's instructions to generate a series of nested deletions from both clones. The resulting truncated

10 inserts were then sequenced from the M13 forward primer present on the pWKS30 vector with the "Taq Dye Deoxy Terminator Cycle Sequencing Kit" using an Applied Biosystems Inc. (Foster City, CA) automated sequencer model 373A according to the manufacturer's

15 recommendations.

B. Sequence Analysis

After the insert was sequenced in both directions, the nucleotide sequence revealed an open reading frame

20 consisting of 525 nucleotides (including the stop codon) encoding a protein composed of 174 amino acid residues having a predicted molecular weight of 18,000 Daltons and a pI of 9.93. The nucleotide and deduced amino acid sequences are presented in Figure 1 (SEQ ID NO:1; SEQ ID

25 NO:2).

To confirm the correct expression of the cloned gene, the N-terminal amino acid sequence of the native 22 kDa surface protein derived from *Neisseria meningitidis* strain 608B was determined in order to compare it with the amino

30 acid sequence deduced from the nucleotide sequencing data.

Outer membrane preparation derived from *Neisseria meningitidis* strain 608B was resolved by electrophoresis on a 14% SDS-PAGE gel and transferred onto a polyvinylidene difluoride membrane (Millipore Products,

35 Bedford MA) according to a previously described method

[Sambrook et al., Molecular Cloning; a laboratory manual, Cold Spring Harbor Laboratory Press (1989)]. The 22 kDa protein band was excised from the gel and then subjected to Edman degradation using the Applied Biosystems Inc. 5 (Foster City, CA) model 473A automated protein sequencer following the manufacturer's recommendations. The amino acid sequence E-G-A-S-G-F-Y-V-Q-A corresponded to amino acids 1-10 (SEQ ID NO:2) of the open reading frame, indicating that the *Neisseria meningitidis* strain 608B, 22 10 kDa surface protein has a 19 amino acid leader peptide (amino acid residues -19 to -1 of SEQ ID NO:2).

A search of established databases confirmed that the *Neisseria meningitidis* strain 608B, 22 kDa surface protein (SEQ ID NO:2) or its gene (SEQ ID NO:1) have not been 15 described previously.

C. High Yield Expression And Purification of The Recombinant *Neisseria meningitidis* 22 kDa Surface Protein

20 The following process was developed in order to maximize the production and purification of the recombinant *Neisseria meningitidis* 22 kDa surface protein expressed in *Escherichia coli*. This process is based on the observation that the recombinant 22 kDa surface 25 protein produced by *Escherichia coli* strain BL21(DE3) [Studier and Moffat, J. Mol. Biol., 189, p. 113 (1986)] carrying the plasmid pNP2202 can be found in large amounts in the outer membrane, but can also be obtained from the culture supernatant in which it is the most abundant 30 protein. The culture supernatant was therefore the material used to purify the recombinant 22 kDa protein using affinity chromatography (Figure 6A).

To generate an affinity chromatography matrix, monoclonal antibodies Me-2, Me-3 and Me-5 (described in 35 Example 2) were immobilized on CNBr-activated sepharose 4B

(Pharmacia Biotech Inc., Piscataway, NJ) according to the manufacturer's instructions.

To prepare the culture supernatant, an overnight culture of *Escherichia coli* strain BL21(DE3), harboring
5 the plasmid pNP2202 was inoculated in LB broth (Gibco Laboratories, Grand Island, N.Y.) containing 25 mg/ml of ampicillin (Sigma) and was incubated 4 hours at 37°C with agitation. The bacterial cells were removed from the culture media by two centrifugations at 10,000 Xg for 10
10 minutes at 4°C. The culture supernatant was filtered onto a 0.22 µm membrane (Millipore, Bedford, MA) and then concentrated approximately 100 times using an ultra-filtration membrane (Amicon Co., Beverly, MA) with a molecular cut off of 10,000 Daltons. To completely
15 solubilize the membrane vesicles, Empigen BB (Calbiochem Co., LaJolla, CA) was added to the concentrated culture supernatant to a final concentration of 1% (vol/vol). The suspension was incubated at room temperature for one hour, dialyzed overnight against several liters of 10 mM Tris-
20 HCl buffer, pH 7.3 containing 0.05% Empigen BB(vol/vol) and centrifuged at 10,000Xg for 20 minutes at 4°C. The antigen preparation was added to the affinity matrix and incubated overnight at 4°C with constant agitation. The gel slurry was poured into a chromatography column and
25 washed extensively with 10 mM Tris-HCl buffer, pH 7.3 containing 0.05% Empigen BB (vol/vol). The recombinant 22 kDa protein was then eluted from the column with 1 M LiCl in 10 mM Tris-HCl buffer, pH 7.3. The solution containing the eluted protein was dialyzed extensively against
30 several liters of 10 mM Tris-HCl buffer, pH 7.3 containing 0.05% Empigen BB. Coomassie Blue and silver stained SDS-Page gels [Tsai and Frasch, Analytical Biochem., 119, pp. 19 (1982)] were used to evaluate the purity of the recombinant 22 kDa surface protein at each step of the
35 purification process and representative results are

presented in Figure 6A. Silver staining of the gels clearly demonstrated that the purification process generated a fairly pure recombinant 22 kDa protein with only a very small quantity of *Escherichia coli*

5 lipopolysaccharide.

The resistance to proteolytic cleavage of the purified recombinant 22 kDa surface protein was also verified and the results are presented in Figure 6B. Purified recombinant 22 kDa surface protein was treated as
10 described in Example 1 with α -chymotrypsin and trypsin at 2 mg per mg of protein and with 2 IU of proteinase K per mg of protein for 1 hour at 37°C with constant shaking. No reduction in the amount of protein was observed after any of these treatments. In comparison, partial or
15 complete digestion depending on the enzyme selected was observed for the control protein which was in this case bovine serum albumin (BSA, Sigma). Furthermore, longer periods of treatment did not result in any modification of the protein. These latter results demonstrated that
20 transformed *Escherichia coli* cells can express the complete recombinant 22 kDa surface protein and that this protein is also highly resistant to the action of these three proteolytic enzymes as was the native protein found in *Neisseria meningitidis*. In addition, the purified
25 recombinant 22 kDa surface protein which is not embedded in the outer membrane of *Escherichia coli* is still highly resistant to the action of the proteolytic enzymes.

We also verified the effect of the enzymatic treatments on the antigenic properties of the recombinant
30 22 kDa protein. As determined by ELISA and Western immunoblotting, the monoclonal antibodies described in Example 2 readily recognized the recombinant 22 kDa surface protein that was purified according to the process described above (Figure 6C). Moreover, the reactivity of
35 monoclonal antibody Me-5, as well as the reactivity of

other 22 kDa protein-specific monoclonal antibodies, with the purified recombinant 22 kDa surface protein was not altered by any of the enzyme treatments, thus confirming that the antigenic properties of the recombinant 22 kDa protein seem similar to the ones described for the native protein.

Important data were presented in Example 3 and can be summarized as follows:

- 1) the complete nucleotide and amino acid sequences of the *Neisseria meningitidis* 22 kDa surface protein were obtained (SEQ ID NO:1; SEQ ID NO:2);
- 2) N-terminal sequencing of the native protein confirmed that the *Neisseria meningitidis* 22 kDa gene was indeed cloned;
- 3) this protein was not described previously;
- 4) it is possible to transform a host such as *Escherichia coli* and obtain expression of the recombinant *Neisseria meningitidis* 22 kDa surface protein in high yield;
- 5) it is possible to obtain the recombinant protein free of other *Neisseria meningitidis* molecules and almost free of components produced by *Escherichia coli*;
- 6) the purified recombinant 22 kDa surface protein remains highly resistant to the action of proteolytic enzymes such as α -chymotrypsin, trypsin and proteinase K; and
- 7) the antigenic properties of the recombinant 22 kDa protein compare to the ones described for the native *Neisseria meningitidis* 22 kDa surface protein.

EXAMPLE 4 Molecular Conservation Of The Gene Coding for the *Neisseria meningitidis* 22 kDa Surface Protein

To verify the molecular conservation among *Neisseria* isolates of the gene coding for the *Neisseria meningitidis* 22 kDa surface protein, a DNA dot blot hybridization assay was used to test different *Neisseria* species and other
5 bacterial species. First, the 525 base pair gene coding for the *Neisseria meningitidis* 22 kDa surface protein was amplified by PCR, purified on agarose gel and labeled by random priming with the non radioactive DIG DNA labeling and detection system (Boehringer Mannheim, Laval, Canada)
10 following the manufacturer's instructions.

The DNA dot blot assay was done according to the manufacturer's instructions (Boehringer Mannheim). Briefly, the bacterial strains to be tested were dotted onto a positively charge nylon membrane (Boehringer
15 Mannheim), dried and then treated as described in the DIG System's user's guide for colony lifts. Pre-hybridizations and hybridizations were done at 42°C with solutions containing 50% formamide (Sigma). The pre-hybridization solution also contained 100 mg/ml of
20 denatured herring sperm DNA (Boehringer Mannheim) as an additional blocking agent to prevent non-specific hybridization of the DNA probe. The stringency washes and detection steps using the chemiluminescent lumigen PPD substrate were also done as described in the DIG System's
25 user's guide.

For the 71 *Neisseria meningitidis* strains tested the results obtained with monoclonal antibody Me-7 and the 525 base pair DNA probe were in perfect agreement. According to the results, all the *Neisseria meningitidis* strains
30 tested have the *Neisseria meningitidis* 22 kDa surface protein gene and they express the protein since they were all recognized by the monoclonal antibody, thus confirming that this protein is highly conserved among the *Neisseria meningitidis* isolates (Table 2).

The DNA probe also detected the gene coding for the *Neisseria meningitidis* 22 kDa surface protein in all *Neisseria gonorrhoeae* strains tested.

On the contrary, the monoclonal antibody Me-7 reacted
5 only with 2 out of the 16 *Neisseria gonorrhoeae* strains tested indicating that the specific epitope is somehow absent, inaccessible or modified in *Neisseria gonorrhoeae* strains, or that most of the *Neisseria gonorrhoeae* strains do not express the protein even if they have the coding
10 sequence in their genome (Table 2).

A good correlation between the two detection methods was also observed for *Neisseria lactamica*, since only one strain of *Neisseria lactamica* was found to have the gene without expressing the protein (Table 2). This result
15 could also be explained by the same reasons presented in the last paragraph.

This may indicate that, although the 22kDa is not expressed, or not accessible on the surface of *Neisseria gonorrhoeae* strains, the 22kDa protein-coding gene of the
20 *Neisseria gonorrhoeae* and *Neisseria lactamica* strains may be used for construction of recombinant plasmids used for the production of the 22kDa surface protein or analogs. All such protein or analogs may be used for the prevention, detection, or diagnosis of *Neisseria*
25 infections. More particularly, such infections may be selected from infections from *Neisseria meningitidis*, *Neisseria gonorrhoeae*, and *Neisseria lactamica*. Therefore, the 22kDa surface protein or analogs, may be used for the manufacture of a vaccine against such infections.
30 Moreover, the 22kDa protein or analogs, may be used for the manufacture of a kit for the detection or diagnosis of such infections.

The results obtained with *Moraxella catharralis* strains showed that out of the 5 strains tested, 3 reacted
35 with monoclonal antibody Me-7, but none of them reacted

with the DNA probe indicating that the gene coding for the *Neisseria meningitidis* 22 kDa surface protein is absent from the genome of these strains (Table 2).

Several other *Neisserial* species as well as other
 5 bacterial species (see footnote, Table 2) were tested and none of them were found to be positive by any of the two tests. This latter result seems to indicate that the gene for the 22 kDa surface protein is shared only among closely related species of *Neisseriaceae*.

10

Table 2. Reactivity of the 525 base pair DNA probe and monoclonal antibody Me-7 with different *Neisseria* species

<i>Neisseria</i> species (number of strains tested) ¹	Number of strains identified by	
	Monoclonal antibody Me-7	DNA probe
<i>Neisseria meningitidis</i> (71)	71	71
<i>Moraxella catharallis</i> (5)	3	0
<i>Neisseria gonorrhoeae</i> (16)	2	16
<i>Neisseria lactamica</i> (5)	4	5

15 The following *Neisserial* species and other bacterial species were also tested with the two assays and gave negative results: 1 *Neisseria cinerea*, 1 *Neisseria flava*, 1 *Neisseria flavescens*, 2 *Neisseria mucosa*, 4 *Neisseria perflava/sicca*, 1 *Neisseria perflava*, 1 *N. sicca*, 1 *N. subflava*, 1 *Alcaligenes faecalis* (ATCC 8750), 1 *Bordetella pertussis* (9340), 1 *Bordetella bronchiseptica*, 1 *Citrobacter freundii* (ATCC 2080), 1 *Edwardsiella tarda* (ATCC 15947), 1
 20 *Enterobacter cloaca* (ATCC 23355), 1 *Enterobacter aerogenes* (ATCC 13048), 1 *Escherichia coli*, 1 *Flavobacterium odoratum*, 1 *Haemophilus influenzae* type b (Eagan strain), 1 *Klebsiella pneumoniae* (ATCC 13883), 1 *Proteus rettgeri* (ATCC 25932), 1 *Proteus vulgaris* (ATCC 13315), 1 *Pseudomonas aeruginosa* (ATCC 9027), 1 *Salmonella typhimurium* (ATCC 14028), 1 *Serratia marcescens* (ATCC 8100), 1 *Shigella flexneri* (ATCC 12022), 1 *Shigella sonnei*
 25 (ATCC 9290), and 1 *Xanthomonas maltophilia*.

In conclusion, the DNA hybridization assay clearly indicated that the gene coding for the *Neisseria*

meningitidis 22 kDa surface protein is highly conserved among the pathogenic *Neisseria*. Furthermore, the results obtained clearly showed that this DNA probe could become a valuable tool for the rapid and direct detection of pathogenic *Neisseria* bacteria in clinical specimen. This probe could even be refined to discriminate between the *Neisseria meningitidis* and *Neisseria gonorrhoeae*.

EXAMPLE 5 Bacteriolytic And Protective Properties Of The Monoclonal Antibodies

The bacteriolytic activity of the purified *Neisseria meningitidis* 22 kDa surface protein-specific monoclonal antibodies was evaluated in vitro according to a method described previously [Brodeur et al., Infect. Immun., 50, p. 510 (1985); Martin et al., Infect. Immun., 60, p. 2718 (1992)]. In the presence of a guinea pig serum complement, purified monoclonal antibodies Me-1 and Me-7 efficiently killed *Neisseria meningitidis* strain 608B. Relatively low concentrations of each of these monoclonal antibodies reduced by more than 50% the number of viable bacteria. The utilization of higher concentrations of purified monoclonal antibodies Me-1 and Me-7 resulted in a sharp decrease (up to 99%) in the number of bacterial colony forming units. Importantly, the bacteriolytic activity of these monoclonal antibodies is complement dependent, since heat-inactivation of the guinea pig serum for 30 minutes at 56°C completely abolished the killing activity. The other monoclonal antibodies did not exhibit significant bacteriolytic activity against the same strain. The combined, representative results of several experiments are presented in Figure 7, wherein the results shown for Me-7 are representative and consistent with the results obtained for Me-1. The results shown for Me-2 are

representative and consistent with the results obtained for the other monoclonal antibodies Me-3, Me-5 and Me-6.

A mouse model of infection, which was described previously by one of the inventors [Brodeur et al, Infect. Immun., 50, p. 510 (1985); Brodeur et al., Can. J. Microbiol., 32, p. 33 (1986)] was used to assess the protective activity of each monoclonal antibody. Briefly, Balb/c mice were injected intraperitoneally with 600 μ l of ascitic fluid containing the monoclonal antibodies 18 hours before the bacterial challenge. The mice were then challenged with one ml of a suspension containing 1000 colony forming units of *Neisseria meningitidis* strain 608B, 4% mucin (Sigma) and 1.6% hemoglobin (Sigma). The combined results of several experiments are presented in Table 3. It is important to note that only the bacteriolytic monoclonal antibodies Me-1 and Me-7 protected the mice against experimental *Neisseria meningitidis* infection. Indeed, the injection of ascitic fluid containing these two monoclonal antibodies before the bacterial challenge significantly increased the rate of survival of Balb/c mice to 70% or more compared to the 9% observed in the control groups receiving either 600 μ l Sp2/0 induced ascitic fluid or 600 μ l ascitic fluid containing unrelated monoclonal antibodies. Results have also indicated that 80% of the mice survived the infection if they were previously injected with 400 μ g of protein A purified Me-7 18 hours before the bacterial challenge. Subsequent experiments are presently being done to determine the minimal antibody concentration necessary to protect 50% of the mice. Lower survival rates from 20 to 40% were observed for the other *Neisseria meningitidis* 22 kDa surface protein-specific monoclonal antibodies.

Table 3. Evaluation of the immunoprotective potential of the 22 kDa surface protein-specific monoclonal antibodies against *Neisseria meningitidis* strain 608B (B:2a:P1.2)

5

Monoclonal antibodies	Number of living mice after challenge		% of survival
	24 h	72 h	
Me-1	29/30	23/30	76
Me-2	17/20	3/20	25
Me-3	5/10	2/10	20
Me-5	11/20	8/20	40
Me-7	10/10	7/10	70
purified Me-7	13/15	12/15	80
Control	31/100	9/100	9

In conclusion, the results clearly indicated that an antibody specific for the *Neisseria meningitidis* 22 kDa surface protein can efficiently protect mice against an experimental lethal challenge. The induction of protective antibodies by an antigen is one of the most important criteria to justify further research on potential vaccine candidate.

15 **EXAMPLE 6 Immunization With Purified Recombinant 22 kDa Surface Protein Confers Protection Against Subsequent Bacterial Challenge**

Purified recombinant 22 kDa surface protein was prepared according to the protocol presented in Example 3, and was used to immunize Balb/c mice to determine its protective effect against challenge with a lethal dose of

Neisseria meningitidis 608B (B:2a:P1.2). It was decided to use the purified recombinant protein instead of the native meningococcal protein in order to insure that there was no other meningococcal antigen in the vaccine preparation used during these experiments. The mouse model of infection used in these experiments was described previously by one of the inventors [Brodeur et al., Infect. Immun., 50, p. 510 (1985); Brodeur et al., Can. J. Microbiol., 32, p. 33 (1986)]. The mice were each injected subcutaneously three times at three-week intervals with 100 µl of the antigen preparation containing either 10 or 20 µg per mouse of the purified recombinant 22 kDa surface protein. QuilA was the adjuvant used for these experiments at a concentration of 25 µg per injection. Mice in the control groups were injected following the same procedure with either 10 or 20 µg of BSA, 20 µg of concentrated culture supernatant of *Escherichia coli* strain BL21(DE3) carrying the plasmid pWKS30 without the insert gene for the meningococcal protein prepared as described in Example 3, or phosphate-buffered saline. Serum samples from each mouse were obtained before each injection in order to analyze the development of the immune response against the recombinant protein. Two weeks following the third immunization the mice in all groups were injected intraperitoneally with 1 ml of a suspension containing 1000 colony forming units of *Neisseria meningitidis* strain 608B in 4% mucin (Sigma) and 1.6% hemoglobin (Sigma).

The results of these experiments are presented in Table 4. Eighty percent (80%) of the mice immunized with the purified recombinant 22 kDa surface protein survived the bacterial challenge compared to 0 to 42% in the control groups. Importantly, the mice in the control group injected with concentrated *Escherichia coli* culture supernatant were not protected against the bacterial

challenge. This latter result clearly demonstrated that the components present in the culture media and the *Escherichia coli*'s antigens that might be present in small amounts after purification do not contribute to the
 5 observed protection against *Neisseria meningitidis*.

Table 4. Immunization With Purified Recombinant 22 kDa Surface Protein Confers Protection Against Subsequent Bacterial Challenge with *Neisseria meningitidis* 608B (B:2a:P1.2) strain.

10

Experiment	Group	Number of living mice after challenge			% of survival
		24 h	48 h	72 h	
1	10 µg of purified 22kDa	20/20		16/20	80
	10 µg of BSA	17/19		8/19	42
2	20 µg of purified 22 kDa protein	9/10	8/10	8/10	80
	20 µg of concentrated <i>E. coli</i> supernatant	7/10	5/10	2/10	20
	20 µg of BSA	6/10	4/10	2/10	20
	Phosphate buffered saline	8/10	0/10	0/10	0

CONCLUSION

The injection of purified recombinant 22 kDa surface
 15 protein greatly protected the immunized mice against the development of a lethal infection by *Neisseria meningitidis*.

Antibodies according to this invention are exemplified by murine hybridoma cell lines producing
 20 monoclonal antibodies Me-1 and Me-7 deposited in the American Type Culture Collection in Rockville, Maryland,

USA on July 21, 1995. The deposits were assigned accession numbers BB 11959 (Me-1) and BB 11958 (Me-7).

EXAMPLE 7 Sequence analysis of other strains of
5 *Neisseria meningitidis* and of *Neisseria*
 gonorrhoeae

The 2.75 kb *clai*I digested DNA fragment containing the gene coding for the 22kDa surface protein was isolated
10 from the genomic DNA of the different strains of *Neisseria meningitidis* and *Neisseria gonorrhoeae* as described in Example 3.

a) MCH88 strain: The nucleotide sequence of strain MCH88
15 (clinical isolate) is presented in Figure 8 (SEQ ID NO:3). From experimental evidence obtained from strain 608B (Example 3), a putative leader sequence was deduced corresponding to amino acid -19 to -1 (M-K-K-A-L-A-A-L-I-A-L-A-L-P-A-A-A-L-A). A search of established databases
20 confirmed that 22kDa surface protein from *Neisseria meningitidis* strain MCH 188 (SEQ ID NO:4) or its gene (SEQ ID NO:3) have not been described previously.

b) Z4063 strain: The nucleotide sequence of strain Z4063
25 (Wang J.-F. et al. Infect. Immun., 60, p.5267 (1992)) is presented in Figure 9 (SEQ ID NO:5). From experimental evidence obtained from strain 608B (Example 3), a putative leader sequence was deduced corresponding to amino acid -19 to -1 (M-K-K-A-L-A-T-L-I-A-L-A-L-P-A-A-A-L-A). A search
30 of established databases confirmed that 22kDa surface protein from *Neisseria meningitidis* strain Z4063 (SEQ ID NO:6) or its gene (SEQ ID NO:5) have not been described previously.

c) *Neisseria gonorrhoeae* strain b2: The nucleotide sequence of *Neisseria gonorrhoeae* strain b2 (serotype 1, Nat.Ref. Center for *Neisseria*, LCDC, Ottawa, Canada) is described in Figure 10 (SEQ ID NO:7). From experimental
5 evidence obtained from strain 608B (Example 3), a putative leader sequence was deduced corresponding to amino acid -19 to -1 (M-K-K-A-L-A-A-L-I-A-L-A-L-P-A-A-A-L-A). A search of established databases confirmed that 22kDa surface protein from *Neisseria gonorrhoeae* strain b2 (SEQ ID NO:8)
10 or its gene (SEQ ID NO:7) have not been described previously.

Figure 11 shows the consensus sequence established from the DNA sequence of all four strains tested. The MCH88 strain showed an insertion of one codon (TCA) at
15 nucleotide 217, but in general the four strains showed striking homology.

Figure 12 depicts the homology between the deduced amino acid sequence obtained from the four strains. There is greater than 90% identity between all four strains.
20

Example 8 Immunological response of rabbits and monkeys to the 22kDa *Neisseria meningitidis* surface protein

Rabbits and monkeys were immunized with the
25 recombinant 22kDa protein to assess the antibody response in species other than the mouse.

a) Rabbits

Male New Zealand rabbits were immunized with outer membrane preparations obtained from *E. coli* strain JM109
30 with the plasmid pN2202 or with the control plasmid pWKS30 (the strain and the plasmids are described in Example 3). The lithium chloride extraction used to obtain these outer membrane preparations was performed in a manner previously

described by the inventors [Brodeur et al, Infect. Immun. 50, 510 (1985)]. The protein content of these preparations were determined by the Lowry method adapted to membrane fractions [Lowry et al, J. Biol. Chem. 193, 265 (1951)]. The rabbits were injected subcutaneously and intramuscularly at several sites twice at three week intervals with 150 µg of one of the outer membrane preparations described above. QuilA, at a final concentration of 20% (vol./vol.) (CedarLane Laboratories, Hornby, Ont., Canada), was the adjuvant used for these immunizations. The development of the specific humoral response was analyzed by ELISA using outer membrane preparations extracted from *Neisseria meningitidis* strain 608B (B:2a:P1.2) as coating antigen and by Western immunoblotting following methods already described by the inventors [Brodeur et al., Infect. Immun. 50, 510 (1985); Martin et al, Eur. J. Immunol. 18, 601 (1988)]. Alkaline phosphatase or peroxylase-labeled Donkey anti-rabbit immunoglobulins (Jackson ImmunoResearch Laboratories, West Grove, PA) were used for these assays.

The injection of *E. coli* outer membrane preparation containing the 22 kDa recombinant protein in combination with QuilA adjuvant induced in the rabbit a strong specific humoral response of 1/32,000 as determined by ELISA (Figure 13). The antibodies induced after the injection of the recombinant 22 kDa protein reacted with the purified recombinant 22 kDa protein, but more importantly they also recognized the native protein as

expressed, folded and embedded in the outer membrane of *Neisseria meningitidis*. Western Immunoblotting experiments clearly indicated that the antibodies present after the second injection recognized on nitrocellulose membrane the same protein band as the one revealed by Mab Me-2 (described in Example 2), which is specific for the 22 kDa protein.

b) Monkeys

Two *Macaca fascicularis* (cynomolgus) monkeys were respectively immunized with two injections of 100 µg (K28) and 200 µg (I276) of affinity purified recombinant 22 kDa protein per injection. The methods used to produce and purify the protein from *E. coli* strain BL21De3 were described in Example 3. Alhydrogel, at a final concentration of 20% (vol./vol.) (CedarLane Laboratories, Hornby, Ont., Canada), was the adjuvant used for these immunizations. The monkeys received two intramuscular injections at three weeks interval. A control monkey (K65) was immunized with an unrelated recombinant protein preparation following the same procedures. The sera were analyzed as described above. Alkaline phosphatase or Peroxydase-labeled Goat anti-human immunoglobulins (Jackson ImmunoResearch Laboratories, West Grove, PA) were used for these assays.

The specific antibody response of monkey K28 which was immunized with 100µg of purified protein per injection appeared faster and was stronger than the one observed for

monkey I276 which was injected with 200µg of protein (Figure 14). Antibodies specific for the native 22 kDa protein as detected by Western immunoblotting were already present in the sera of the immunized monkeys twenty one
5 days after the first injection, but were absent in the sera of the control monkey after two injections of the control antigen.

Conclusion

The data presented in Examples 2 and 5 clearly showed
10 that the injection of the recombinant 22 kDa protein can induce a protective humoral response in mice which is directed against *Neisseria meningitidis* strains. More importantly, the results presented in this example demonstrate that this immunological response is not
15 restricted to only one species, but this recombinant surface protein can also stimulate the immune system of other species such as rabbit or monkey.

Example 9 Epitope mapping of the 22kDa *Neisseria meningitidis* protein

20

Neisseria meningitidis 22 kDa surface protein was epitope mapped using a method described by one of the inventors [Martin et al. Infect. Immun. (1991): 59:1457-
25 1464]. Identification of the linear epitopes was accomplished using 18 overlapping synthetic peptides covering the entire *Neisseria meningitidis* 22 kDa protein sequence derived from strain 608B (Figure 15) and

hyperimmune sera obtained after immunization with this protein. The identification of immunodominant portions on the 22 kDa protein may be helpful in the design of new efficient vaccines. Furthermore, the localization of these
5 B-cell epitopes also provides valuable information about the structural configuration of the protein in the outer membrane of *Neisseria meningitidis*.

All peptides were synthesized by BioChem Immunosystems Inc. (Montreal, Canada) with the Applied Biosystems
10 (Foster City, Calif.) automated peptide synthesizer. Synthetic peptides were purified by reverse-phase high-pressure liquid chromatography. Peptides CS-845, CS-847, CS-848, CS-851, CS-852 and CS-856 (Figure 15) were solubilized in a small volume of 6M guanidine-HCl (J.T.
15 Baker, Ontario, Canada) or dimethyl sulfoxide (J.T. Baker). These peptides were then adjusted to 1 mg/ml with distilled water. All the other peptides were freely soluble in distilled water and were also adjusted to 1 mg/ml.

20 Peptide enzyme-linked immunosorbent assays (ELISA) were performed by coating synthetic peptides onto microtitration plates (Immulon 4, Dynatech Laboratories Inc., Chantilly, VA) at a concentration of 50 µg/ml in 50 mM carbonate buffer, pH 9.6. After overnight incubation
25 at room temperature, the plates were washed with phosphate-buffered saline (PBS) containing 0.05% (wt/vol) Tween 20 (Sigma Chemical Co., St.-Louis, Mo.) and blocked with PBS containing 0.5% (wt/vol) bovine serum albumin

- (Sigma). Sera obtained from mice and monkeys immunized with affinity purified recombinant 22 kDa surface protein were diluted and 100 μ l per well of each dilution were added to the ELISA plates and incubated for 1 h at 37°C.
- 5 The plates were washed three times, and 100 μ l of alkaline phosphatase-conjugated goat anti-mouse or anti-human immunoglobulins (Jackson ImmunoResearch Laboratories, West Grove, PA) diluted according to the manufacturer's recommendations was added. After incubation for 1 h at
- 10 37°C, the plates were washed and 100 μ l of diethanolamine (10% (vol/vol), pH 9.8) containing p-nitro-phenylphosphate (Sigma) at 1 mg/ml was added. After 60 min., the reaction (λ =410 nm) was read spectrophotometrically with a microplate reader.
- 15 Mouse and monkey antisera obtained after immunization with affinity purified recombinant 22 kDa protein (Example 8) were successfully used in combination with eighteen overlapping synthetic peptides to localize B-cell epitopes on the protein. These epitopes are clustered within three
- 20 antigenic domains on the protein.

The first region is located between amino acid residues 51 and 86. Computer analysis using different algorithms suggested that this region has the highest probability of being immunologically important since it is

25 hydrophilic and surface exposed. Furthermore, comparison of the four protein sequences which is presented in Figure 12 indicates that one of the major variation, which is the

insertion of one amino acid residue at position 73, is also located in this region.

The antisera identified a second antigenic domain located between amino acid residues 110 and 140.

5 Interestingly, the sequence analysis revealed that seven out of the fourteen amino acid residues that are not conserved among the four protein sequences are clustered within this region of the protein.

A third antigenic domain located in a highly
10 conserved portion of the protein, between amino acid residues 31 and 55, was recognized only by the monkeys' sera.

Example 10 Heat-inducible expression vector for the
15 **large scale production of the 22 kDa**
 surface protein

The gene coding for the *Neisseria meningitidis* 22 kDa surface protein was inserted into the plasmid p629 [George et al. Bio/technology 5: 600-603 (1987)]. A
20 cassette of the bacteriophage λ cI857 temperature sensitive repressor gene, from which the functional Pr promoter has been deleted, is carried by the plasmid p629 that uses the PL promoter to control the synthesis of the 22kDa surface protein. The inactivation of the cI857
25 repressor by a temperature shift from 30°C to temperatures above 38°C results in the production of the protein encoded by the plasmid. The induction of gene expression in *E. coli* cells by a temperature shift is advantageous

for large scale fermentation since it can easily be achieved with modern fermentors. Other inducible expression vectors usually require the addition of specific molecules like lactose or isopropylthio- β -D-galactoside (IPTG) in the culture media in order to induce the expression of the desired gene.

A 540 nucleotide fragment was amplified by PCR from the *Neisseria meningitidis* strain 608B genomic DNA using the following two oligonucleotide primers (OCRR8: 5'-
10 TAATAGATCTATGAAAAAGCACTTGCCAC-3' and OCRR9: 3'-CACGCGCAGTTTAAAGACTTCTAGATTA-5'). These primers correspond to the nucleotide sequences found at both ends of the 22 kDa gene. To simplify the cloning of the PCR product, a Bgl II (AGATCT) restriction site was incorporated into the
15 nucleotide sequence of these primers. The PCR product was purified on agarose gel before being digested with Bgl II.

This Bgl II fragment of approximately 525 base pairs was then inserted into the Bgl II and Bam HI sites of the plasmid p629. The plasmid containing the PCR product
20 insert named pNP2204 was used to transform *E. coli* strain DH5 α F' IQ. A partial map of the plasmid pNP2204 is presented in Figure 16. The resulting colonies were screened with *Neisseria meningitidis* 22 kDa surface-protein specific monoclonal antibodies described in
25 Example 2. Western blot analysis of the resulting clones clearly indicated that the protein synthesized by *E. coli* was complete and migrated on SDS-PAGE gel like the native *Neisseria meningitidis* 22 kDa surface protein. Plasmid

DNA was purified from the selected clone and then sequenced. The nucleotide sequence of the insert present in the plasmid perfectly matched the nucleotide sequence of the gene coding for the *Neisseria meningitidis* 22 kDa protein presented in Figure 1.

To study the level of synthesis of the 22 kDa surface protein, the temperature-inducible plasmid pNP2204 was used to transform the following *E. coli* strains: W3110, JM105, BL21, TOPP1, TOPP2 and TOPP3. The level of synthesis of the 22 kDa surface protein and the localization of the protein in the different cellular fractions were determined for each strain. Shake flask cultures in LB broth (Gibco BRL, Life Technologies, Grand Island, NY) indicated that a temperature shift from 30°C to 39°C efficiently induced the expression of the gene. Time course evaluation of the level of synthesis indicated that the protein appeared, as determined on SDS-PAGE gel, as soon as 30 min after induction and that the amount of protein increased constantly during the induction period.

Expression levels between 8 to 10 mg of 22 kDa protein per liter were determined for *E. coli* strains W3110 and TOPP1. For both strains, the majority of the 22 kDa protein is incorporated in the bacterial outer membrane.

**Example 11 Purification of the *Neisseria meningitidis*
22kDa protein**

Since the vast majority of the 22 kDa protein is found embedded in the outer membrane of *E. coli* strains, the purification protocol presented in this Example is different from the one already described in Example 3 where a large amount of protein was released in the culture supernatant. An overnight culture incubated at 30°C of either *E. coli* strain W3110 or TOPP1 harboring the plasmid pNP2204 was inoculated in LB broth containing 50 µg/ml of Ampicillin (Sigma) and was grown at 30°C with agitation (250 rpm) until it reached a cell density of 0.6 ($\lambda=600\text{nm}$), at which point the incubation temperature was shifted to 39°C for three to five hours to induce the production of the protein. The bacterial cells were harvested by centrifugation at 8,000 xg for 15 minutes at 4°C and washed twice in phosphate buffered saline (PBS), pH 7.3. The bacterial cells were ultrasonically broken (ballistic disintegration or mechanical disintegration with a French press may also be used). Unbroken cells were removed by centrifugation at 5,000 xg for 5 minutes and discarded. The outer membranes were separated from cytoplasmic components by centrifugation at 100,000 xg for 1 h at 10°C. The membrane-containing pellets were resuspended in a small volume of PBS, pH 7.3. To solubilize the 22 kDa surface protein from the membranes, detergents such as Empigen BB (Calbiochem Co., LaJolla, CA), Zwittergent-3,14 (Calbiochem Co.), or β -

octylglucoside (Sigma) were used. The detergent was added to the membrane fraction at final concentration of 3% and the mixture was incubated for 1 h at 20°C. The non soluble material was removed by centrifugation at 100,000
5 xg for 1 h at 10°C.

The 22 kDa protein was efficiently solubilized by either three of the detergents, however β -octylglucoside had the advantage of easily removing several unwanted membrane proteins since they were not solubilized and
10 could be separated from the supernatant by centrifugation.

To remove the detergent, the 22 kDa containing supernatant was dialyzed extensively against several changes of PBS buffer. Proteinase K treatment (as in Example 1) can be used to further remove unwanted proteins
15 from the 22kDa surface protein preparation. Differential precipitation using ammonium sulfate or organic solvents, and ultrafiltration are two additional steps that can be used to remove unwanted nucleic acid and lipopolysaccharide contaminants from the proteins before
20 gel permeation and ion-exchange chromatography can be efficiently used to obtain the purified 22 kDa protein. Affinity chromatography, as described in Example 3, can also be used to purify the 22 kDa protein.

25 **Example 12 Use of 22kDa surface protein As a Human Vaccine**

To formulate a vaccine for human use, appropriate 22kDa surface protein antigens may be selected from the

polypeptides described herein. For example, one of skill in the art could design a vaccine around the 22kDa polypeptide or fragments thereof containing an immunogenic epitope. The use of molecular biology techniques is particularly well-suited for the preparation of substantially pure recombinant antigens.

The vaccine composition may take a variety of forms. These include, for example, solid, semi-solid, and liquid dosage forms, such as powders, liquid solutions or suspensions, and liposomes. Based on our belief that the 22kDa surface protein antigens of this invention may elicit a protective immune response when administered to a human, the compositions of this invention will be similar to those used for immunizing humans with other proteins and polypeptides, e.g. tetanus and diphtheria. Therefore, the compositions of this invention will preferably comprise a pharmaceutically acceptable adjuvant such as incomplete Freund's adjuvant, aluminum hydroxide, a muramyl peptide, a water-in-oil emulsion, a liposome, an ISCOM or CTB, or a non-toxic B subunit form cholera toxin. Most preferably, the compositions will include a water-in-oil emulsion or aluminum hydroxide as adjuvant.

The composition would be administered to the patient in any of a number of pharmaceutically acceptable forms including intramuscular, intradermal, subcutaneous or topic. Preferably, the vaccine will be administered intramuscularly.

Generally, the dosage will consist of an initial injection, most probably with adjuvant, of about 0.01 to 10 mg, and preferably 0.1 to 1.0 mg of 22kDa surface protein antigen per patient, followed most probably by one or more
5 booster injections. Preferably, boosters will be administered at about 1 and 6 months after the initial injection.

A consideration relating to vaccine development is the question of mucosal immunity. The ideal mucosal
10 vaccine will be safely taken orally or intranasally as one or a few doses and would elicit protective antibodies on the appropriate surfaces along with systemic immunity. The mucosal vaccine composition may include adjuvants, inert particulate carriers or recombinant live vectors.

15 The anti-22kDa surface protein antibodies of this invention are useful for passive immunotherapy and immunoprophylaxis of humans infected with *Neisseria meningitidis* or related bacteria such as *Neisseria gonorrhoeae* or *Neisseria lactamica*. The dosage forms and
20 regimens for such passive immunization would be similar to those of other passive immunotherapies.

An antibody according to this invention is exemplified by a hybridoma producing MAbs Me-1 or Me-7 deposited in the American Type Culture Collection in
25 Rockville, Maryland, USA on July 21, 1995, and identified as Murine Hybridoma Cell Lines, Me-1 and Me-7 respectively. These deposits were assigned accession numbers HB 11959 (Me-1) and HB 11958 (Me-7).

While we have described herein a number of
embodiments of this invention, it is apparent that our
basic embodiments may be altered to provide other
embodiments that utilize the compositions and processes of
5 this invention. Therefore, it will be appreciated that the
scope of this invention includes all alternative
embodiments and variations that are defined in the
foregoing specification and by the claims appended
thereto; and the invention is not to be limited by the
10 specific embodiments which have been presented herein by
way of example.

SEQUENCE LISTING

- 5 (1) GENERAL INFORMATION:
- (i) APPLICANT: Brodeur, Bernard R
Martin, Denis
Hamel, Josee
10 Rioux, Clement
- (ii) TITLE OF INVENTION: PROTEINASE K RESISTANT SURFACE PROTEIN
OF NEISSERIA MENINGITIDIS
- 15 (iii) NUMBER OF SEQUENCES: 26
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Goudreau Gage Dubuc & Martineau Walker
(B) STREET: 800 Place Victoria, Suite 3400, Tour de la
20 Bourse
(C) CITY: Montreal
(D) STATE: Quebec
(E) COUNTRY: Canada
(F) ZIP: H4Z 1E9
- 25 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
30 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
35 (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 08/406,362
(B) FILING DATE: 17-MAR-1995
40
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US (PROVIS)60/001,983
(B) FILING DATE: 04-AUG-1995
- 45 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Leclerc/Dubuc/Prince, Alain/Jean/Gaetan
(C) REFERENCE/DOCKET NUMBER: BIOVAC-1 PCT
- (ix) TELECOMMUNICATION INFORMATION:
50 (A) TELEPHONE: 514-397-7400
(B) TELEFAX: 514-397-4382
- (2) INFORMATION FOR SEQ ID NO:1:
- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 830 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
60 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: *Neisseria meningitidis*
 (B) STRAIN: 608B

10 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 143..667

(ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 15 (B) LOCATION: 143..199

(ix) FEATURE:
 (A) NAME/KEY: mat_peptide
 20 (B) LOCATION: 200..667

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

25 TCGGCAAGC AGCCGGATAC CGCTACGTAT CTTGAAGTAT TGAAAATATT ACGATGCAAA 60
AAAGAAAATT TAAGTATAAT ACAGCAGGAT TCTTTAACGG ATTCTTAACA ATTTTCTTAA 120
CTGACCATAA AGGAACCAAAA AT ATG AAA AAA GCA CTT GCC ACA CTG ATT GCC 172
Met Lys Lys Ala Leu Ala Thr Leu Ile Ala
30 -19 -15 -10
CTC GCT CTC CCG GCC GCC GCA CTG GCG GAA GGC GCA TCC GGC TTT TAC 220
Leu Ala Leu Pro Ala Ala Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr
-5 1 5
35 GTC CAA GCC GAT GCC GCA CAC GCA AAA GCC TCA AGC TCT TTA GGT TCT 268
Val Gln Ala Asp Ala Ala His Ala Lys Ala Ser Ser Ser Leu Gly Ser
10 15 20
40 GCC AAA GGC TTC AGC CCG CGC ATC TCC GCA GGC TAC CGC ATC AAC GAC 316
Ala Lys Gly Phe Ser Pro Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp
25 30 35
45 CTC CGC TTC GCC GTC GAT TAC ACG CGC TAC AAA AAC TAT AAA GCC CCA 364
Leu Arg Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro
40 45 50 55
50 TCC ACC GAT TTC AAA CTT TAC AGC ATC GGC GCG TCC GCC ATT TAC GAC 412
Ser Thr Asp Phe Lys Leu Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp
60 65 70
55 TTC GAC ACC CAA TCG CCC GTC AAA CCG TAT CTC GGC GCG CGC TTG AGC 460
Phe Asp Thr Gln Ser Pro Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser
75 80 85
CTC AAC CGC GCC TCC GTC GAC TTG GGC GGC AGC GAC AGC TTC AGC CAA 508
Leu Asn Arg Ala Ser Val Asp Leu Gly Gly Ser Asp Ser Phe Ser Gln
90 95 100
60 ACC TCC ATC GGC CTC GGC GTA TTG ACG GGC GTA AGC TAT GCC GTT ACC 556
Thr Ser Ile Gly Leu Gly Val Leu Thr Gly Val Ser Tyr Ala Val Thr
105 110 115

```

CCG AAT GTC GAT TTG GAT GCC GGC TAC CGC TAC AAC TAC ATC GGC AAA 604
 Pro Asn Val Asp Leu Asp Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys
 120 125 130 135

5 GTC AAC ACT GTC AAA AAC GTC CGT TCC GGC GAA CTG TCC GTC GGC GTG 652
 Val Asn Thr Val Lys Asn Val Arg Ser Gly Glu Leu Ser Val Gly Val
 140 145 150

10 CGC GTC AAA TTC TGATATGCGC CTTATTCTGC AAACCGCCGA GCCTTCGGCG 704
 Arg Val Lys Phe
 155

GTTTTGTTTT CTGCCACCGC AACTACACAA GCCGGCGGTT TTGTACGATA ATCCCGAATG 764

15 CTGCGGCTTC TGCCGCCCTA TTTTTTGAGG AATCCGAAT GTCCAAAACC ATCATCCACA
 824
 ACA 830

20 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 174 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30 Met Lys Lys Ala Leu Ala Thr Leu Ile Ala Leu Ala Leu Pro Ala Ala
 -19 -15 -10 -5

35 Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala
 1 5 10

His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro
 15 20 25

40 Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp
 30 35 40 45

Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu
 50 55 60

45 Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser Pro
 65 70 75

50 Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Val
 80 85 90

Asp Leu Gly Gly Ser Asp Ser Phe Ser Gln Thr Ser Ile Gly Leu Gly
 95 100 105

55 Val Leu Thr Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp
 110 115 120 125

Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys Val Asn Thr Val Lys Asn
 130 135 140

60 Val Arg Ser Gly Glu Leu Ser Val Gly Val Arg Val Lys Phe
 145 150 155

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 5 (A) LENGTH: 710 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 15 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: MCH88
- (ix) FEATURE:
 20 (A) NAME/KEY: CDS
 (B) LOCATION: 116..643
- (ix) FEATURE:
 25 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 116..172
- (ix) FEATURE:
 30 (A) NAME/KEY: mat_peptide
 (B) LOCATION: 173..643

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

35 GTATCTTGAG GCATTGAAAA TATTACAATG CAAAAAGAAA ATTTTCAGTAT AATACGGCAG 60
   GATTCTTTAA CGGATTCTTA ACCATTTTTC TCCCTGACCA TAAAGGAATC AAGAT ATG 118
                                     Met
                                     -19
40
   AAA AAA GCA CTT GCC GCA CTG ATT GCC CTC GCC CTC CCG GCC GCC GCA 166
   Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala Ala
   -15 -10 -5
45 CTG GCG GAA GGC GCA TCC GGC TTT TAC GTC CAA GCC GAT GCC GCA CAC 214
   Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala His
   1 5 10
50 GCC AAA GCC TCA AGC TCT TTA GGT TCT GCC AAA GGC TTC AGC CCG CGC 262
   Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro Arg
   15 20 25 30
   ATC TCC GCA GGC TAC CGC ATC AAC GAC CTC CGC TTC GCC GTC GAT TAC 310
   Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp Tyr
   55 35 40 45
   ACG CGC TAC AAA AAC TAT AAA CAA GTC CCA TCC ACC GAT TTC AAA CTT 358
   Thr Arg Tyr Lys Asn Tyr Lys Gln Val Pro Ser Thr Asp Phe Lys Leu
   50 55 60
60
   TAC AGC ATC GGC GCG TCC GCC ATT TAC GAC TTC GAC ACC CAA TCC CCC 406
   Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser Pro
   65 70 75

```

```

      GTC AAA CCG TAT CTC GGC GCG CGC TTG AGC CTC AAC CGC GCC TCC GTC 454
      Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Val
          80                      85                      90
5      GAC TTT AAC GGC AGC GAC AGC TTC AGC CAA ACC TCC ACC GGC CTC GGC 502
      Asp Phe Asn Gly Ser Asp Ser Phe Ser Gln Thr Ser Thr Gly Leu Gly
          95                      100                      105                      110
10     GTA TTG GCG GGC GTA AGC TAT GCC GTT ACC CCG AAT GTC GAT TTG GAT 550
      Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp
                      115                      120                      125
15     GCC GGC TAC CGC TAC AAC TAC ATC GGC AAA GTC AAC ACT GTC AAA AAT 598
      Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys Val Asn Thr Val Lys Asn
                      130                      135                      140
      GTC CGT TCC GGC GAA CTG TCC GCC GGC GTA CGC GTC AAA TTC TGATATACGC 650
      Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe
          145                      150                      155
20     GTTATTCGCG AAACCGCCGA GCCTTTCGGC GGTTTTGTTT TCCGCGCGCG CAACTACACA 710

```

25

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 175 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala
-19          -15          -10          -5
40  Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala
          1          5          10
      His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro
          15          20          25
45  Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp
      30          35          40          45
      Tyr Thr Arg Tyr Lys Asn Tyr Lys Gln Val Pro Ser Thr Asp Phe Lys
50          50          55          60
      Leu Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser
          65          70          75
55  Pro Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser
          80          85          90
      Val Asp Phe Asn Gly Ser Asp Ser Phe Ser Gln Thr Ser Thr Gly Leu
          95          100          105
60  Gly Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu
      110          115          120          125

```

(2) INFORMATION FOR SEQ ID NO:5:

40	CACCCATCCG CCGCGTGTATG CCGCCACCAC CATTTAAGG CAACGCGCGG GTTAAACGGCT	60
	TTGCCGTCGG CAAAGCAGCC GGATACCGCT ACGTATCTTG AAGTATTAAT AATATTACGA	120
45	TGCAAAAAGA AAATTTAAGT ATAATAAAGC AGAATTCTTT AACGGATTCT TAACAATTTT	180
	TCTAACTGAC CATAAAGGAA CCAAAT ATG AAA AAA GCA CTT GCC ACA CTG	231
	Met Lys Lys Ala Leu Ala Thr Leu	
	-19 -15	
50		
	ATT GCC CTC GCT CTC CCG GCC GCC GCA CTG GCG GAA GGC GCA TCC GGC	279
	Ile Ala Leu Ala Leu Pro Ala Ala Ala Leu Ala Glu Gly Ala Ser Gly	
	-10 -5 1 5	
55	TTT TAC GTC CAA GCC GAT GCC GCA CAC GCA AAA GCC TCA AGC TCT TTA	327
	Phe Tyr Val Gln Ala Asp Ala Ala His Ala Lys Ala Ser Ser Ser Leu	
	10 15 20	
60	GGT TCT GCC AAA GGC TTC AGC CCG CGC ATC TCC GCA GGC TAC CGC ATC	375
	Gly Ser Ala Lys Gly Phe Ser Pro Arg Ile Ser Ala Gly Tyr Arg Ile	
	25 30 35	


```

AAC GAC CTC CGC TTC GCC GTC GAT TAC ACG CGC TAC AAA AAC TAT AAA 423
Asn Asp Leu Arg Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys
      40                      45                      50

5  GCC CCA TCC ACC GAT TTC AAA CTT TAC AGC ATC GGC GCG TCC GCC ATT 471
   Ala Pro Ser Thr Asp Phe Lys Leu Tyr Ser Ile Gly Ala Ser Ala Ile
      55                      60                      65

10 TAC GAC TTC GAC ACC CAA TCG CCC GTC AAA CCG TAT CTC GGC GCG CGC 519
   Tyr Asp Phe Asp Thr Gln Ser Pro Val Lys Pro Tyr Leu Gly Ala Arg
      70                      75                      80                      85

15 TTG AGC CTC AAC CGC GCC TCC GTC GAC TTG GGC GGC AGC GAC AGC TTC 567
   Leu Ser Leu Asn Arg Ala Ser Val Asp Leu Gly Gly Ser Asp Ser Phe
      90                      95                      100

   AGC CAA ACC TCC ACC GGC CTC GGC GTA TTG GCG GGC GTA AGC TAT GCC 615
   Ser Gln Thr Ser Thr Gly Leu Gly Val Leu Ala Gly Val Ser Tyr Ala
      105                      110                      115

20 GTT ACC CCG AAT GTC GAT TTG GAT GCC GGC TAC CGC TAC AAC TAC ATC 663
   Val Thr Pro Asn Val Asp Leu Asp Ala Gly Tyr Arg Tyr Asn Tyr Ile
      120                      125                      130

25 GGC AAA GTC AAC ACT GTC AAA AAC GTC CGT TCC GGC GAA CTG TCC GCC 711
   Gly Lys Val Asn Thr Val Lys Asn Val Arg Ser Gly Glu Leu Ser Ala
      135                      140                      145

30 GGT GTG CGC GTC AAA TTC TGATATGCGC CTTATTCTGC AAACCGCCGA 759
   Gly Val Arg Val Lys Phe
      150                      155

GCCTTCGGCG GTTTTGTTTT CTGCCACCGC AACTACACAA GCCGGCGGTT TTGTACGATA 819

35 ATCCCGAATG CTGCGGCTTC TGCCGCCCTA T 850

```

(2) INFORMATION FOR SEQ ID NO:6:

```

40 (i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 174 amino acids
    (B) TYPE: amino acid
    (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

50 Met Lys Lys Ala Leu Ala Thr Leu Ile Ala Leu Ala Leu Pro Ala Ala
   -19                      -15                      -10                      -5

   Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala
      1                      5                      10

55 His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro
   15                      20                      25

   Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp
   30                      35                      40                      45

60 Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu
      50                      55                      60

```

```

      Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser Pro
                65                      70                      75
5  Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Val
      80                      85                      90
      Asp Leu Gly Gly Ser Asp Ser Phe Ser Gln Thr Ser Thr Gly Leu Gly
      95                      100                      105
10 Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp
    110                      115                      120
      Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys Val Asn Thr Val Lys Asn
      130                      135                      140
15 Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe
      145                      150                      155

```

20 (i) INFORMATION FOR SEQ ID NO:7:

```

      (i) SEQUENCE CHARACTERISTICS:
          (A) LENGTH: 810 base pairs
          (B) TYPE: nucleic acid
25      (C) STRANDEDNESS: double
          (D) TOPOLOGY: linear

      (ii) MOLECULE TYPE: DNA (genomic)

30      (iii) HYPOTHETICAL: NO

          (iv) ANTI-SENSE: NO

          (vi) ORIGINAL SOURCE:
35      (A) ORGANISM: Neisseria gonorrhoeae
          (B) STRAIN: b2

          (ix) FEATURE:
40      (A) NAME/KEY: CDS
          (B) LOCATION: 241..765

          (ix) FEATURE:
          (A) NAME/KEY: sig_peptide
45      (B) LOCATION: 241..297

          (ix) FEATURE:
          (A) NAME/KEY: mat_peptide
          (B) LOCATION: 298..765

```

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

CCCCGCCTTT GCGGTTTTTTT CCAAACCGTT TGCAAGTTTC ACCCATCCGC CGCGTGATGC 60
55 CGCCGTTTAA GGGCAACGCG CGGGTTAACG GATTTCGCGT CGGCAAAGCA GCCGGATGCC 120
      GCGCGTATC TTGAGGCATT GAAAATATTA CGATGCAAAA AGAAAATTTT AGTATAATAC 180
      GGCAGGATTC TTTAACGGAT TATTAACAAT TTTTCTCCCT GACCATAAAG GAACCAAAAT 240
60 ATG AAA AAA GCA CTT GCC GCA CTG ATT GCC CTC GCA CTC CCG GCC GCC 288
Met Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala
-19          -15          -10          -5

```

5 GCA CTG GCG GAA GGC GCA TCC GGC TTT TAC GTC CAA GCC GAT GCC GCA 336
 Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala
 1 5 10
 10 CAC GCC AAA GCC TCA AGC TCT TTA GGT TCT GCC AAA GGC TTC AGC CCG 384
 His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro
 15 20 25
 15 CGC ATC TCC GCA GGC TAC CGC ATC AAC GAC CTC CGC TTC GCC GTC GAT 432
 Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp
 30 35 40 45
 15 TAC ACG CGC TAC AAA AAC TAT AAA GCC CCA TCC ACC GAT TTC AAA CTT 480
 Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu
 50 55 60
 20 TAC AGC ATC GGC GCG TCC GTC ATT TAC GAC TTC GAC ACC CAA TCG CCC 528
 Tyr Ser Ile Gly Ala Ser Val Ile Tyr Asp Phe Asp Thr Gln Ser Pro
 65 70 75
 25 GTC AAA CCG TAT TTC GGC GCG CGC TTG AGC CTC AAC CGC GCT TCC GCC 576
 Val Lys Pro Tyr Phe Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Ala
 80 85 90
 25 CAC TTG GGC GGC AGC GAC AGC TTC AGC AAA ACC TCC GCC GGC CTC GGC 624
 His Leu Gly Gly Ser Asp Ser Phe Ser Lys Thr Ser Ala Gly Leu Gly
 95 100 105
 30 GTA TTG GCG GGC GTA AGC TAT GCC GTT ACC CCG AAT GTC GAT TTG GAT 672
 Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp
 110 115 120 125
 35 GCC GGC TAC CGC TAC AAC TAC GTC GGC AAA GTC AAC ACT GTC AAA AAC 720
 Ala Gly Tyr Arg Tyr Asn Tyr Val Gly Lys Val Asn Thr Val Lys Asn
 130 135 140
 40 GTC CGT TCC GGC GAA CTG TCC GCC GGC GTG CGC GTC AAA TTC TGATATACGC 772
 Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe
 145 150 155
 GTTATTCCGC AAACCGCCGA GCCTTCGGCG GTTTTTTG 810

45 (2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 174 amino acids

(B) TYPE: amino acid

50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

55 Met Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala
 -19 -15 -10 -5
 60 Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala
 1 5 10
 His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro
 15 20 25

Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp
 30 35 40 45
 5 Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu
 50 55 60
 Tyr Ser Ile Gly Ala Ser Val Ile Tyr Asp Phe Asp Thr Gln Ser Pro
 65 70 75
 10 Val Lys Pro Tyr Phe Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Ala
 80 85 90
 15 His Leu Gly Gly Ser Asp Ser Phe Ser Lys Thr Ser Ala Gly Leu Gly
 95 100 105
 Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp
 110 115 120 125
 20 Ala Gly Tyr Arg Tyr Asn Tyr Val Gly Lys Val Asn Thr Val Lys Asn
 130 135 140
 Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe
 145 150 155
 25

(2) INFORMATION FOR SEQ ID NO:9:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: protein
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: 608B
 40
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
 45 Met Lys Lys Ala Leu Ala Thr Leu Ile Ala Leu Ala Leu Pro Ala Ala
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:10:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 55 (ii) MOLECULE TYPE: protein
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: 608B
 60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

5 Leu Ala Leu Pro Ala Ala Ala Leu Ala Glu Gly Ala Ser Gly Phe
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

15 (vi) ORIGINAL SOURCE:

 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: 608B

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

 Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala His Ala Lys
 1 5 10 15

25 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35 (vi) ORIGINAL SOURCE:

 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: 608B

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

 Ala Ala His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly
 1 5 10 15

45 (2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

50 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

55 (vi) ORIGINAL SOURCE:

 (A) ORGANISM: Neisseria meningitidis
 (B) STRAIN: 608B

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly Ser Ala Lys Gly Phe Ser Pro Arg Ile Ser Ala Gly Tyr Arg
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp Tyr
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu Tyr Ser Ile Gly Ala
1 5 10 15

5

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

15

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Neisseria meningitidis

(B) STRAIN: 608B

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser
1 5 10 15

25

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Neisseria meningitidis

(B) STRAIN: 608B

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Phe Asp Thr Gln Ser Pro Val Lys Pro Tyr Leu Gly Ala Arg Leu
1 5 10 15

45

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

50

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

55

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Neisseria meningitidis

(B) STRAIN: 608B

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Val Asp Leu Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ser Val Asp Leu Gly Gly Ser Asp Ser Phe Ser Gln Thr Ser Ile
1 5 10 15

20 (2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

30 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ser Gln Thr Ser Ile Gly Leu Gly Val Leu Thr Gly Val Ser Tyr
1 5 10 15

40 (2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

50 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Thr Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp Ala
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Neisseria meningitidis*

(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Val	Asp	Leu	Asp	Ala	Gly	Tyr	Arg	Tyr	Asn	Tyr	Ile	Gly	Lys	Val
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Neisseria meningitidis*

(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Tyr	Ile	Gly	Lys	Val	Asn	Thr	Val	Lys	Asn	Val	Arg	Ser	Gly	Glu
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Neisseria meningitidis*

(B) STRAIN: 608B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Val	Arg	Ser	Gly	Glu	Leu	Ser	Val	Gly	Val	Arg	Val	Lys	Phe
1				5					10				

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 25 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Neisseria meningitidis
(B) STRAIN: 608B

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

20 Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr
1 5 10 15
Asp Phe Lys Leu Tyr Ser Ile Gly Ala
20 25

We claim:

1. An isolated antigen or fragment thereof which is immunologically accessible on greater than 50% of known strains of *Neisseria meningitidis*.
5
2. The isolated antigen or fragment of claim 1 which is immunologically accessible on about 99% of known strains of *Neisseria meningitidis*.
10
3. The isolated antigen or fragment of claim 1 in which immunological accessibility is determined by using an agglutination assay, an ELISA, a RIA, an immunoblotting assay, a dot-enzyme assay, a surface accessibility assay, or a combination of these assays.
15
4. The isolated antigen or fragment of claim 1 which is a protein.
- 20 5. The protein of claim 4 which has a molecular weight of about 22 kilodaltons.
6. The protein of claim 4 which has a molecular weight of about 18 kilodaltons.
25
7. The protein of claim 5 which has the amino acid sequence selected from the sequences of: Figure 1 (SEQ ID NO:2), Figure 8 (SEQ ID NO:4), Figure 9 (SEQ ID NO:6), and Figure 10 (SEQ ID NO:8).
30
8. The protein of claim 5 which has the amino acid sequence of Figure 1 (SEQ ID NO:2).
9. The protein of claim 5 in substantially pure form.

10. The protein of claim 9, wherein said substantially pure form is obtained by the steps of:
- a) isolating a culture of *Neisseria meningitidis* bacteria,
 - b) isolating an outer membrane portion from the culture of the bacteria; and
 - c) isolating said antigen from the outer membrane portion.
11. The protein of claim 10, wherein step (c) includes the additional step of treating the outer membrane portion with proteinase K followed by protein fractionation.
12. A DNA sequence encoding at least a portion of at least one antigen of the *Neisseria meningitidis* 22 kDa surface protein, said sequence being selected from the group consisting of:
- a) the DNA sequence of Figure 1 (SEQ ID NO:1);
 - b) the DNA sequence of Figure 8 (SEQ ID NO:3);
 - c) the DNA sequence of Figure 9 (SEQ ID NO:5);
 - d) the DNA sequence of Figure 10 (SEQ ID NO:7);
 - e) analogues or derivatives of the foregoing DNA sequences;
 - f) DNA sequences degenerate to any of the foregoing DNA sequences; and
 - g) fragments of any of the foregoing DNA sequences;
- wherein said sequences encode a product that displays the immunological activity of the *Neisseria meningitidis* 22 kDa surface protein.
13. A DNA sequence encoding at least a portion of at least one antigen of the *Neisseria meningitidis* 22 kDa

surface protein, said sequence being selected from the group consisting of:

- a) the DNA sequence of Figure 1 (SEQ ID NO:1);
 - b) analogues or derivatives of the foregoing DNA
5 sequence;
 - c) DNA sequences degenerate to any of the foregoing DNA sequences; and
 - d) fragments of any of the foregoing DNA sequences;
- wherein said sequences encode a product that displays
10 the immunological activity of the *Neisseria meningitidis* 22 kDa surface protein.

14. A DNA sequence according to claim 12, wherein said analog is selected from the DNA of *Neisseria*
15 *gonorrhoeae*.

15. A DNA sequence according to claim 12, wherein said analog is selected from the DNA of *Neisseria*
20 *lactamica*.

16. A DNA sequence of the formula of Figure 1 (SEQ ID NO:1) from base 143 to base 667.

17. A DNA sequence of the formula of Figure 1 (SEQ ID
25 NO:1) from base 200 to base 667.

18. A DNA sequence of the formula of Figure 8 (SEQ ID NO:3) from base 116 to base 643.

19. A DNA sequence of the formula of Figure 8 (SEQ ID
30 NO:3) from base 173 to base 643.

20. A DNA sequence of the formula of Figure 9 (SEQ ID NO:5) from base 208 to base 732.

21. A DNA sequence of the formula of Figure 9 (SEQ ID NO:5) from base 265 to base 732.
- 5 22. A DNA sequence of the formula of Figure 10 (SEQ ID NO:7) from base 241 to base 765.
23. A DNA sequence of the formula of Figure 10 (SEQ ID NO:7) from base 298 to base 765.
- 10 24. A fragment of a DNA sequence according to claim 13, wherein said fragment is selected from the group consisting of: one of the peptides illustrated in Figure 15 (SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, and SEQ ID NO:26).
- 15 25. A fragment of a DNA sequence comprising from amino acid 31 to amino acid 55 of Figure 1 (SEQ ID NO:1).
- 20 26. A fragment of a DNA sequence comprising from amino acid 51 to amino acid 86 of Figure 1 (SEQ ID NO:1).
- 25 27. A fragment of a DNA sequence comprising from amino acid 110 to amino acid 140 of Figure 1 (SEQ ID NO:1).
- 30 28. A recombinant DNA molecule comprising a DNA sequence selected from the group consisting of the DNA sequences of claims 13, 16, or 17, and one or more expression control sequences operatively linked to the DNA sequence.

29. A recombinant DNA molecule comprising a DNA sequence selected from the group consisting of the DNA sequences of claims 14, 15, or any one of claims 18 to 27, and one or more expression control sequences operatively linked to the DNA sequence.
30. The recombinant DNA molecule of claim 28, wherein said expression control sequence is an inducible expression vector.
31. The recombinant DNA molecule of claim 29, wherein said expression control sequence is an inducible expression vector.
32. The recombinant DNA molecule of claim 30 or 31, wherein said vector is induced by stimuli selected from: temperature, presence of lactose, and presence of IPTG.
33. The recombinant DNA molecule of claim 32, wherein said vector is selected from: λ PL, λ PR, TAC, T7, T3, LAC, and TRP promoters.
34. A plasmid selected from the group consisting of: pNP2202, pNP2203, and pNP2204.
35. A unicellular host transformed with a recombinant DNA molecule of claim 28.
36. A unicellular host transformed with a recombinant DNA molecule of claim 29.

37. The unicellular host according to claim 35 or 36,
wherein the host is selected from the group consisting
of: strains of *E.coli* JM109, *E.coli* BL21(DE3), *E.coli*
DH5 α F' IQ, *E.coli* W3110, *E.coli* JM105, *E.coli* BL21,
5 *E.coli* TOPP1, *E.coli* TOPP2, and *E.coli* TOPP3.
38. The unicellular host according to claim 35 or 36,
wherein the host is selected from the group consisting
of strains of *E.coli* JM109 and *E.coli* BL21(DE3).
- 10 39. A *Neisseria meningitidis* 22kDa surface protein in
substantially pure form obtained by the culturing of a
unicellular host according to claim 35 or 36 and
isolating said protein.
- 15 40. A polypeptide coded for by a DNA sequence of any one
of claims 13, 16 and 17.
41. A polypeptide coded for by a DNA sequence of any one
20 of claims 14, 15 and 18 to 27.
42. A method for producing a DNA sequence comprising the
steps of culturing the unicellular host of claim 35 or
36 and isolating said DNA sequence.
- 25 43. A method for producing a polypeptide comprising the
steps of culturing the unicellular host of claim 35 or
36 and isolating said polypeptide.
- 30 44. A polypeptide in substantially pure form as obtained
by the method of claim 43.

45. A method for isolating the antigen of claim 1 comprising:
- a) isolating a culture of *Neisseria meningitidis* bacteria,
 - 5 b) isolating an outer membrane portion from the culture of the bacteria; and
 - c) isolating said antigen from the outer membrane portion.
- 10 46. The method of claim 45, wherein step (c) includes the additional steps of treating the outer membrane portion with proteinase K followed by protein fractionation.
- 15 47. A pharmaceutical composition comprising one or more antigens or fragments thereof according to any one of claims 1 to 5, 8, 13, 16, and 17.
- 20 48. A pharmaceutical composition comprising one or more antigens or fragments thereof according to claim 40.
49. A pharmaceutical composition comprising one or more antigens or fragments thereof according to claim 41.
- 25 50. A pharmaceutical composition comprising one or more antigens or fragments thereof according to any one of claims 6, 7, 9, 12, 14, 15, and 18 to 27.
- 30 51. The pharmaceutical composition of claim 49, which is a vaccine.
52. The pharmaceutical composition of claim 50, which is a vaccine.

53. The pharmaceutical composition of claim 49, further comprising one or more pharmaceutically acceptable excipients.
- 5 54. The pharmaceutical composition of claim 50, further comprising one or more pharmaceutically acceptable excipients.
- 10 55. A method for preventing infection of a patient by *Neisseria meningitidis* comprising the administration of a pharmaceutically effective amount of the vaccine of claim 51 or 52.
- 15 56. The use of a pharmaceutically effective amount of the *Neisseria meningitidis* 22kDa surface protein or a fragment, analogue, or derivative thereof for the prevention of *Neisseria meningitidis* infection in humans.
- 20 57. The use of a pharmaceutically effective amount of the *Neisseria meningitidis* 22kDa surface protein or a fragment, analogue, or derivative thereof for the prevention of *Neisseria gonorrhoeae* infection in humans.
- 25 58. The use of the *Neisseria meningitidis* 22 kDa surface protein or a fragment, analogue, or derivative thereof for the manufacture of a vaccine for the prevention of *Neisseria meningitidis* infection in humans.
- 30 59. The use of the *Neisseria meningitidis* 22 kDa surface protein or a fragment, analogue, or derivative thereof for the manufacture of a vaccine for the prevention of *Neisseria gonorrhoeae* infection in humans.

60. An antibody or fragment thereof that specifically binds to a protein with a molecular weight of approximately 22 kilodaltons present on greater than 50% of known strains of *Neisseria meningitidis*.
61. The antibody or fragment of claim 60 that specifically binds to about 99% of known strains of *Neisseria meningitidis*.
62. The antibody or fragment of claim 60 which is a monoclonal antibody or fragment thereof.
63. The monoclonal antibody or fragment of claim 62 which is of murine origin.
64. The monoclonal antibody or fragment of claim 63 which is of an IgG isotype.
65. The monoclonal antibody or fragment of claim 62 which is Me-1, Me-2, Me-3, Me-5, Me-6, or Me-7.
66. The monoclonal antibody of claim 62 which is the monoclonal antibody Me-1 or Me-7.
67. A method for isolating the antibody of claim 60 comprising:
a) introducing a preparation of *Neisseria meningitidis* into a mammal; and
b) isolating serum from the mammal containing said antibody.
68. A method for isolating the monoclonal antibody of claim 62 comprising:

- a) introducing a preparation of *Neisseria meningitidis* to antibody producing cells of a mammal;
- b) fusing the antibody producing cells with myeloma cells to form hybridoma cells; and
- 5 c) isolating said monoclonal antibody from the hybridoma cells.

69. A pharmaceutical composition comprising one or more antibodies or fragments thereof according to any one
10 of claims 60-66.

70. The pharmaceutical composition of claim 69 which is a vaccine.

15 71. The pharmaceutical composition of claim 69, further comprising a pharmaceutically acceptable excipient.

72. The pharmaceutical composition of claim 69, wherein the antibody is Me-1 or Me-7.

20

73. A method for treating a patient infected with or suspected of being infected with *Neisseria meningitidis* comprising the administration of a pharmaceutically effective amount of the vaccine of
25 claim 70.

74. A method for the detection of *Neisseria meningitidis* antigen in a biological sample containing or suspected of containing *Neisseria meningitidis* antigen comprising:
30

- a) isolating the biological sample from a patient;
- b) incubating the antibody or fragment of claim 60 with the biological sample to form a mixture; and

c) detecting specifically bound antibody or bound fragment in the mixture which indicates the presence of *Neisseria meningitidis* antigen.

5 75. The method of claim 72 wherein the antibody is Me-1 or Me-7.

76. A method for the detection of antibody specific to *Neisseria meningitidis* antigen in a biological sample
10 containing or suspected of containing said antibody comprising:

- a) isolating the biological sample from a patient;
- b) incubating the antigen or fragment of claim 1 with the biological sample to form a mixture; and
- 15 c) detecting specifically bound antigen or bound fragment in the mixture which indicates the presence of antibody specific to *Neisseria meningitidis* antigen.

20 77. The method of claim 76 wherein the antigen is the *Neisseria meningitidis* 22 kDa surface protein.

78. A method for the detection of pathogenic *Neisseria* bacteria in a biological sample containing or
25 suspected of containing such bacteria comprising:

- a) isolating the biological sample from a patient;
- b) incubating a DNA probe having the DNA sequence of claim 13 with the biological sample to form a mixture; and
- 30 c) detecting specifically bound DNA probe in the mixture which indicates the presence of *Neisseria* bacteria.

79. A method for the detection of pathogenic *Neisseria* bacteria in a biological sample containing or suspected of containing such bacteria comprising:
- a) isolating the biological sample from a patient;
 - 5 b) incubating a DNA probe having the DNA sequence of claim 12 with the biological sample to form a mixture; and
 - c) detecting specifically bound DNA probe in the mixture which indicates the presence of *Neisseria* bacteria.
- 10
80. The method of claim 78 wherein the DNA probe has the 525 base pair sequence of Figure 1 (SEQ ID NO:1).
81. The method of claim 78 wherein the DNA probe has a portion of the 525 base pair sequence of Figure 1 (SEQ ID NO:1).
- 15
82. The method of claim 79 wherein the DNA probe has the full or a portion of the 528 base pair sequence of Figure 8 (SEQ ID NO:3).
- 20
83. The method of claim 79 wherein the DNA probe has the full or a portion of the 525 base pair sequence of Figure 9 (SEQ ID NO:5).
- 25
84. The method of claim 79 wherein the DNA probe has the a portion or the full 525 base pair sequence of Figure 10 (SEQ ID NO:7).
- 30
85. The method of claim 78 wherein the DNA probe is an oligomer having a sequence complementary to at least about 6 contiguous nucleotides of the *Neisseria*

meningitidis 22 kDa surface protein of Figure 1 (SEQ ID NO:1).

86. The method of claim 79 wherein the DNA probe is an oligomer having a sequence complementary to at least about 6 contiguous nucleotides of the *Neisseria meningitidis* 22 kDa surface protein of sequences selected from: Figure 8 (SEQ ID NO:3), Figure 9 (SEQ ID NO:5), and Figure 10 (SEQ ID NO:7).
87. The method of claim 85 or 86 which further comprises:
- a) providing a set of oligomers which are primers for a polymerase chain reaction method and which flank the target region; and
 - b) amplifying the target region via the polymerase chain reaction method.
88. A method for the detection of *Neisseria meningitidis* in a patient comprising:
- a) labeling the antibody or fragment of claim 60 with a detectable label;
 - b) administering the labeled antibody or labeled fragment to the patient; and
 - c) detecting specifically bound labeled antibody or labeled fragment in the patient which indicates the presence of *Neisseria meningitidis*.
89. The use of a pharmaceutically effective amount of an antibody specific to the *Neisseria meningitidis* 22 kDa surface protein for the prevention of *Neisseria meningitidis* infection in humans.
90. The use of the *Neisseria meningitidis* 22 kDa surface protein or a fragment, analogue, or derivative thereof

for the manufacture of a kit for the detection or diagnosis of *Neisseria meningitidis* infection in humans.

FIGURE 1

	TCGGCAAAGC AGCCGGATAC CGCTACGTAT CTTGAAGTAT TGAAAAATATT ACGATGCAAA	60
5	AAAGAAAATT TAAGTATAAT ACAGCAGGAT TCTTTAACGG ATTCTTAACA ATTTTCTTAA	120
	CTGACCATAA <u>AGGAACCAAA</u> AT ATG AAA AAA GCA CTT GCC ACA CTG ATT GCC	172
	Met Lys Lys Ala Leu Ala Thr Leu Ile Ala	-10
	-19 -15	
10	CTC GCT CTC CCG GCC GCC GCA CTG GCG GAA GGC GCA TCC GGC TTT TAC	220
	Leu Ala Leu Pro Ala Ala Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr	5
	-5 1	
15	GTC CAA GCC GAT GCC GCA CAC GCA AAA GCC TCA AGC TCT TTA GGT TCT	268
	Val Gln Ala Asp Ala Ala His Ala Lys Ala Ser Ser Ser Leu Gly Ser	20
	10 15	
20	GCC AAA GGC TTC AGC CCG CGC ATC TCC GCA GGC TAC CGC ATC AAC GAC	316
	Ala Lys Gly Phe Ser Pro Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp	35
	25 30	
25	CTC CGC TTC GCC GTC GAT TAC ACG CGC TAC AAA AAC TAT AAA GCC CCA	364
	Leu Arg Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro	50 55
	40 45	
30	TCC ACC GAT TTC AAA CTT TAC AGC ATC GGC GCG TCC GCC ATT TAC GAC	412
	Ser Thr Asp Phe Lys Leu Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp	60 65 70
	60	
35	TTC GAC ACC CAA TCG CCC GTC AAA CCG TAT CTC GGC GCG CGC TTG AGC	460
	Phe Asp Thr Gln Ser Pro Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser	75 80 85
	75 80	
40	CTC AAC CGC GCC TCC GTC GAC TTG GGC GGC AGC GAC AGC TTC AGC CAA	508
	Leu Asn Arg Ala Ser Val Asp Leu Gly Gly Ser Asp Ser Phe Ser Gln	90 95 100
	90	
45	ACC TCC ATC GGC CTC GGC GTA TTG ACG GGC GTA AGC TAT GCC GTT ACC	556
	Thr Ser Ile Gly Leu Gly Val Leu Thr Gly Val Ser Tyr Ala Val Thr	105 110 115
	105 110	
50	CGG AAT GTC GAT TTG GAT GCC GGC TAC CGC TAC AAC TAC ATC GGC AAA	604
	Pro Asn Val Asp Leu Asp Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys	120 125 130 135
	120 125	
55	GTC AAC ACT GTC AAA AAC GTC CGT TCC GGC GAA CTG TCC GTC GGC GTG	652
	Val Asn Thr Val Lys Asn Val Arg Ser Gly Glu Leu Ser Val Gly Val	140 145 150
	140 145	
60	CGC GTC AAA TTC TGATATGGCG CTTATTCTGC AAACCGCCGA GCCTTCGGCG	704
	Arg Val Lys Phe	155
	155	
65	GTTTTGTTTT CTGCCACCGC AACTACACAA GCCGGCGGTT TTGTACGATA ATCCCGAATG	764
	CTGCGGCTTC TGCCGCCCTA TTTTGTGAGG AATCCGAAAT GTCCAAAACC ATCATCCACA	824
	CCGACA	830

FIGURE 2




FIGURE 3a




FIGURE 3b




FIGURE 4a




FIGURE 4b




FIGURE 5




FIGURE 6A




FIGURE 6B




FIGURE 6C

NT C T K

NT C T K

FIGURE 7




FIGURE 8

5	GTATCTTGAG GCATTGAAAA TATTACAATG CAAAAAGAAA ATTTTCAGTAT AATACGGCAG	60
	GATTCTTTTAA CGGATTCTTA ACCATTTTTC TCCCTGACCA TAAAGGAATC AAGAT ATG	118
		Met -19
10	AAA AAA GCA CTT GCC GCA CTG ATT GCC CTC GCC CTC CCG GCC GCC GCA	166
	Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala Ala	
	-15 -10 -5	
15	CTG GCG GAA GGC GCA TCC GGC TTT TAC GTC CAA GCC GAT GCC GCA CAC	214
	Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala His	
	1 5 10	
20	GCC AAA GCC TCA AGC TCT TTA GGT TCT GCC AAA GGC TTC AGC CCG CGC	262
	Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro Arg	
	15 20 25 30	
25	ATC TCC GCA GGC TAC CGC ATC AAC GAC CTC CGC TTC GCC GTC GAT TAC	310
	Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp Tyr	
	35 40 45	
30	ACG CGC TAC AAA AAC TAT AAA CAA GTC CCA TCC ACC GAT TTC AAA CTT	358
	Thr Arg Tyr Lys Asn Tyr Lys Gln Val Pro Ser Thr Asp Phe Lys Leu	
	50 55 60	
35	TAC AGC ATC GGC GCG TCC GCC ATT TAC GAC TTC GAC ACC CAA TCC CCC	406
	Tyr Ser Ile Gly Ala Ser Ala Ile Tyr Asp Phe Asp Thr Gln Ser Pro	
	65 70 75	
40	GTC AAA CCG TAT CTC GGC GCG CGC TTG AGC CTC AAC CGC GCC TCC GTC	454
	Val Lys Pro Tyr Leu Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Val	
	80 85 90	
45	GAC TTT AAC GGC AGC GAC AGC TTC AGC CAA ACC TCC ACC GGC CTC GGC	502
	Asp Phe Asn Gly Ser Asp Ser Phe Ser Gln Thr Ser Thr Gly Leu Gly	
	95 100 105 110	
50	GTA TTG CCG GGC GTA AGC TAT GCC GTT ACC CCG AAT GTC GAT TTG GAT	550
	Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp	
	115 120 125	
55	GCC GGC TAC CGC TAC AAC TAC ATC GGC AAA GTC AAC ACT GTC AAA AAT	598
	Ala Gly Tyr Arg Tyr Asn Tyr Ile Gly Lys Val Asn Thr Val Lys Asn	
	130 135 140	
60	GTC CGT TCC GGC GAA CTG TCC GCC GGC GTA CGC GTC AAA TTC TGATATACGC	650
	Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe	
	145 150 155	
65	GTTATTCCGC AAACCGCCGA GCCTTTCGGC GGTTTTGTTC TCCGCCGCCG CAACTACACA	710

FIGURE 9

	CACCCATCCG CCGCGTGATG CCGCCACCAC CATTATAAGG CAACGCGCGG GTTAACGGCT	60
5	TTGCCGTCGG CAAAGCAGCC GGATACCGCT ACGTATCTTG AAGTATTAAA AATATTACGA	120
	TGCAAAAAGA AAATTTAAGT ATAATAAAGC AGAATTCTTT AACGGATTCT TAACAATTTT	180
10	TCTAACTGAC CATAAAGGAA CCAAAAT ATG AAA AAA GCA CTT GCC ACA CTG	231
	Met Lys Lys Ala Leu Ala Thr Leu	
	-19 -15	
15	ATT GCC CTC GGT CTC CCG GCC GCC GCA CTG GCG GAA GGC GCA TCC GGC	279
	Ile Ala Leu Ala Leu Pro Ala Ala Leu Ala Glu Gly Ala Ser Gly	
	-10 -5 1 5	
20	TTT TAC GTC CAA GCC GAT GCC GCA CAC GCA AAA GCC TCA AGC TCT TTA	327
	Phe Tyr Val Gln Ala Asp Ala Ala His Ala Lys Ala Ser Ser Ser Leu	
	10 15 20	
25	GGT TCT GCC AAA GGC TTC AGC CCG CGC ATC TCC GCA GGC TAC CGC ATC	375
	Gly Ser Ala Lys Gly Phe Ser Pro Arg Ile Ser Ala Gly Tyr Arg Ile	
	25 30 35	
30	AAC GAC CTC CGC TTC GCC GTC GAT TAC ACG CGC TAC AAA AAC TAT AAA	423
	Asn Asp Leu Arg Phe Ala Val Asp Tyr Thr Arg Tyr Lys Asn Tyr Lys	
	40 45 50	
35	GCC CCA TCC ACC GAT TTC AAA CTT TAC AGC ATC GGC GCG TCC GCC ATT	471
	Ala Pro Ser Thr Asp Phe Lys Leu Tyr Ser Ile Gly Ala Ser Ala Ile	
	55 60 65	
40	TAC GAC TTC GAC ACC CAA TCG CCC GTC AAA CCG TAT CTC GGC GCG CGC	519
	Tyr Asp Phe Asp Thr Gln Ser Pro Val Lys Pro Tyr Leu Gly Ala Arg	
	70 75 80 85	
45	TTG AGC CTC AAC CGC GCC TCC GTC GAC TTG GGC GGC AGC GAC AGC TTC	567
	Leu Ser Leu Asn Arg Ala Ser Val Asp Leu Gly Gly Ser Asp Ser Phe	
	90 95 100	
50	AGC CAA ACC TCC ACC GGC CTC GGC GTA TTG GCG GGC GTA AGC TAT GCC	615
	Ser Gln Thr Ser Thr Gly Leu Gly Val Leu Ala Gly Val Ser Tyr Ala	
	105 110 115	
55	GTT ACC CCG AAT GTC GAT TTG GAT GCC GGC TAC CGC TAC AAC TAC ATC	663
	Val Thr Pro Asn Val Asp Leu Asp Ala Gly Tyr Arg Tyr Asn Tyr Ile	
	120 125 130	
60	GGC AAA GTC AAC ACT GTC AAA AAC GTC CGT TCC GGC GAA CTG TCC GCC	711
	Gly Lys Val Asn Thr Val Lys Asn Val Arg Ser Gly Glu Leu Ser Ala	
	135 140 145	
65	GGT GTG CGC GTC AAA TTC TGATATGCGC CTTATCTGCG AAACCGCCGA	759
	Gly Val Arg Val Lys Phe	
	150 155	
70	GCCTTCGGCG GTTTTGTTTT CTGCCACCGC AACTACACAA GCCGGCGGTT TTGTACGATA	819
	ATCCCGAATG CTGCGGCTTC TGCCGCCCTA T	850

FIGURE 10

	CCCCGCCTTT GCGGTTTTTTT CCAAACCGTT TGCAAGTTTC ACCCATCCGC CGCGTGATGC	60
5	CGCCGTTTAA GGGCAACGCG CGGGTTAACG GATTGCGCGT CGGCAAAGCA GCCGGATGCC	120
	GCCGCGTATC TTGAGGCATT GAAAATATTA CGATGCAAAA AGAAAATTTT AGTATAATAC	180
10	GGCAGGATTC TTTAACGGAT TATTAACAAT TTTTCTCCCT GACCATAAAG GAACCAAAAT	240
	ATG AAA AAA GCA CTT GCC GCA CTG ATT GCC CTC GCA CTC CCG GCC GCC	288
	Met Lys Lys Ala Leu Ala Ala Leu Ile Ala Leu Ala Leu Pro Ala Ala	
	-19 -15 -10 -5	
15	GCA CTG GCG GAA GGC GCA TCC GGC TTT TAC GTC CAA GCC GAT GCC GCA	336
	Ala Leu Ala Glu Gly Ala Ser Gly Phe Tyr Val Gln Ala Asp Ala Ala	
	1 5 10	
20	CAC GCC AAA GCC TCA AGC TCT TTA GGT TCT GCC AAA GGC TTC AGC CCG	384
	His Ala Lys Ala Ser Ser Ser Leu Gly Ser Ala Lys Gly Phe Ser Pro	
	15 20 25	
25	CGC ATC TCC GCA GGC TAC CGC ATC AAC GAC CTC CGC TTC GCC GTC GAT	432
	Arg Ile Ser Ala Gly Tyr Arg Ile Asn Asp Leu Arg Phe Ala Val Asp	
	30 35 40 45	
	TAC ACG CGC TAC AAA AAC TAT AAA GCC CCA TCC ACC GAT TTC AAA CTT	480
	Tyr Thr Arg Tyr Lys Asn Tyr Lys Ala Pro Ser Thr Asp Phe Lys Leu	
	50 55 60	
30	TAC AGC ATC GGC GCG TCC GTC ATT TAC GAC TTC GAC ACC CAA TCG CCC	528
	Tyr Ser Ile Gly Ala Ser Val Ile Tyr Asp Phe Asp Thr Gln Ser Pro	
	65 70 75	
35	GTC AAA CCG TAT TTC GGC GCG CGC TTG AGC CTC AAC CGC GCT TCC GCC	576
	Val Lys Pro Tyr Phe Gly Ala Arg Leu Ser Leu Asn Arg Ala Ser Ala	
	80 85 90	
40	CAC TTG GGC GGC AGC GAC AGC TTC AGC AAA ACC TCC GCC GGC CTC GGC	624
	His Leu Gly Gly Ser Asp Ser Phe Ser Lys Thr Ser Ala Gly Leu Gly	
	95 100 105	
45	GTA TTG GCG GGC GTA AGC TAT GCC GTT ACC CCG AAT GTC GAT TTG GAT	672
	Val Leu Ala Gly Val Ser Tyr Ala Val Thr Pro Asn Val Asp Leu Asp	
	110 115 120 125	
	GCC GGC TAC CGC TAC AAC TAC GTC GGC AAA GTC AAC ACT GTC AAA AAC	720
	Ala Gly Tyr Arg Tyr Asn Tyr Val Gly Lys Val Asn Thr Val Lys Asn	
	130 135 140	
50	GTC CGT TCC GGC GAA CTG TCC GCC GGC GTG CGC GTC AAA TTC TGATATACGC	772
	Val Arg Ser Gly Glu Leu Ser Ala Gly Val Arg Val Lys Phe	
	145 150 155	
55	GTTATTCGCG AAACCGCCGA GCCTTCGGCG GTTTTTTG	810
60		

FIGURE 11

5	MCH88-ORFG.C.	50	
	608B-ORFA.T.	50	
	Z4063-ORFA.T.	50	
	gono b2-ORFG.A.	50	
	Consensus	ATGAAAAAAG	CACTTGCCRC	ACTGATTGCC	CTCGCHCTCC	CGGCCGCCGC
10	MCH88-ORF	100	
	608B-ORF	100	
	Z4063-ORF	100	
	gono b2-ORF	100	
	Consensus	ACTGGCGGAA	GGCGCATCCG	GCTTTTACGT	CCAAGCCGAT	GCCGCACACG
15	MCH88-ORF	.C.....	150	
	608B-ORF	.A.....	150	
	Z4063-ORF	.A.....	150	
	gono b2-ORF	.C.....	150	
	Consensus	CMAAAGCCTC	AAGCTCTTTA	GGTTCGCCA	AAGGCTTCAG	CCCGCGCATC
20	MCH88-ORF	200	
	608B-ORF	200	
	Z4063-ORF	200	
	gono b2-ORF	200	
	Consensus	TCCGCAGGCT	ACCGCATCAA	CGACCTCCGC	TTCGCCGTGC	ATTACACGCG
25	MCH88-ORF	T.....	250
	608B-ORF	C.....	247
	Z4063-ORF	C.....	247
	gono b2-ORF	C.....	247
	Consensus	CTACAAAAAC	TATAAACAAG	YCCCATCCAC	CGATTTCAAA	CTTTACAGCA
30	MCH88-ORFC.....C..	300
	608B-ORFC.....G..	297
	Z4063-ORFC.....G..	297
	gono b2-ORFT.....G..	298
	Consensus	TGGCGCGGTC	CGYCATTTAC	GACTTCGACA	CCCAATCSCC	CGTCAAACCG
35	MCH88-ORF	...C.....C...T.G	...TAA...	-350
	608B-ORF	...C.....C...T.G	...GGG...	347
	Z4063-ORF	...C.....C...T.G	...GGG...	347
	gono b2-ORF	...T.....T...C.C	...GGG...	347
	Consensus	TATYTCGGCG	CGCGCTTGAG	CCTCAACCGC	GCTTCGGYCS	ACTTKRRCCG
40	MCH88-ORFC...AC...G.....	400
	608B-ORFC...AT...A.....	397
	Z4063-ORFC...AC...G.....	397
	gono b2-ORFA...GC...G.....	397
	Consensus	CAGCGACAGC	TTCAGCMAAA	CCTCCRYCGG	CCTCGGCGTA	TTGRCGGGCG
45	MCH88-ORF	450
	608B-ORF	447
	Z4063-ORF	447
	gono b2-ORF	447
	Consensus	TAAGCTATGC	CGTTACCCCG	AATGTCGATT	TGGATGCCGG	CTACCGCTAC
50	MCH88-ORFA...T.....	500
	608B-ORFA...C.....	497
	Z4063-ORFA...C.....	497
	gono b2-ORFG...C.....	497
	Consensus	AACTACRTCG	GCAAAGTCAA	CACTGTCAAA	AAYGTCCGTT	CCGGCGAACT
55	MCH88-ORFC...C	..A.....	528
	608B-ORFT...C	..G.....	525
	Z4063-ORFC...T	..G.....	525
	gono b2-ORFC...C	..G.....	525
	Consensus	GTCCGYCGGY	GTRCGCGTCA	AATTCTGA

FIGURE 12

5	gonoB2A...	50
	Z4063T...	50
	608BT...	50
	MCH88A...	50
	Consensus	MKKALA.LIA	LALPAAALAE	GASGFYVQAD	AAHAKASSSL	GSAGGFSPRI	50
10	gonoB2V..	99
	Z4063	99
	608B	99
	MCH88QV.....	99
	Consensus	SAGYRINDLR	FAVDYTRYKN	YK-APSTDFK	LYSIGASAIY	DFDTQSPVKP	100
15	gonoB2	.F.....	..AH.....	..K..A....	149
	Z4063T....	149
	608BI....	..T.....	149
	MCH88FN....T....	150
	Consensus	YLGARLSLNR	ASVDLGGSDS	FSQTS.GLGV	LAGVSYAVTP	NVDLDAGYRY	150
20	gonoB2	..V.....	174
	Z4063	174
	608BV..	174
	MCH88	175
	Consensus	NYIGKVNTVK	NVRSGELSAG	VRVKF	175

FIGURE 13




FIGURE 14





FIGURE 15

	MKKALATLIA	LALPAAALAE	GASGFYVQAD	AAHAKASSSL	GSAKGFSPRI	50
5	CS-840		CS-842		CS-844	
		CS-841		CS-843		
10	SAGYRINDLR	FAVDYTRYKN	YKAPSTDFKL	YSIGASAIYD	FDTQSPVKPY	100
		CS-846		CS-848		
	CS-845		CS-847		CS-849	
15		CS-857				
	LGARLSLNRA	SVDLGGSDSF	SQTSIGLGVL	TGVSYAVTPN	VDLDAGYRYN	150
	CS-850		CS-852		CS-854	
20		CS-851		CS-853		
	YIGKVNTVKN	VRSGELSVGV	RVKF			174
25		CS-856				
	CS-855					
30						

FIGURE 16



INTERNATIONAL SEARCH REPORT

International Application No.

PC1/CA 96/00157

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/31 C07K14/22 C07K16/12 C12N15/70 A61K38/16
A61K39/095 G01N33/569 C12Q1/68 C12N1/21 //(C12N1/21,
C12R1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 05703 (GLOBAL TEK INC) 17 March 1994	1-6,9, 10,45, 60-62, 67-79
Y	see the whole document ---	47-52
Y	EP,A,0 273 116 (MAX PLANCK GESELLSCHAFT) 6 July 1988	7,8, 12-14, 16-28, 39-43, 47-52
	see claims; figures --- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

12 August 1996

Date of mailing of the international search report

14. 08. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Delanghe, L

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 96/00157

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MOLECULAR MICROBIOLOGY, vol. 5, no. 8, 1991, pages 1889-1901, XP000578159 K.S.BHAT ET AL.: "The opacity proteins of Neisseria gonorrhoeae strain MS11 are encoded by a family of 11 complete genes." see the whole document ---	7,8, 12-14, 16-28, 39-43
Y	MOLECULAR MICROBIOLOGY, vol. 6, no. 8, 1992, pages 1073-1076, XP000578158 K.S.BHAT ET AL.: "The opacity proteins of Neisseria gonorrhoeae strain MS11 are encoded by a family of 11 complete genes." see the whole document ---	7,8, 12-14, 16-28, 39-43
Y	EMBO JOURNAL, vol. 12, no. 2, 1993, EYNSHAM, OXFORD GB, pages 641-650, XP002010474 EVA-MARIA KUPSCH ET AL.: "Variable opacity (Opa) outer membrane proteins account for the cell tropisms displayed by Neisseria gonorrhoeae for human leukocytes and epithelial cells." see the whole document -----	7,8, 12-14, 16-28, 39-43

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA96/00157

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 55-57, 73, 88, 89
because they relate to subject matter not required to be searched by this Authority, namely:
Although these claims are directed to a method of treatment of the human body as well as diagnostic methods (rule 39.1.(IV) PCT) the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 96/00157

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9405703	17-03-94	AU-B- 5093693 CA-A- 2122630 EP-A- 0625165	29-03-94 17-03-94 23-11-94
EP-A-0273116	06-07-88	NONE	